



LOVE YOUR LIVER AND GET MORE OUT OF LIFE! WITH HEPATIC DETOX



David Woolcott Naturopath

Our liver certainly is a hard worker. Treat it well, live longer and get more out of life. The liver is involved with many different functions of the body. Most of us understand that our liver is the major organ that carries out most of the detoxification, but it does a lot more. It is also responsible for bile production. In fact it produces up to a litre of bile per day. Bile needs to be at the right pH which is somewhere between 7.5 and 8.6. Bile consists mostly of bile acids, water, bile salts, cholesterol, bile pigments and a phospholipid called lecithin, plus several ions. Bile is required for emulsification of fats and for the activation of lipase enzymes needed for digestion.

The liver is the heaviest gland of the body and plays a major role in maintaining proper hormonal balance. It can chemically alter or excrete thyroid hormones and steroid hormones, such as estrogens and aldosterone. The liver is required to carry out detoxification of the bloodstream by degrading various waste materials no longer needed. There are two types of toxins. There are those, which come from outside the human body (exogenous), and those, which are generated by the processes within the body (endogenous). Xenobiotics are chemical compounds which are foreign to the body, (exogenous toxins) such as drugs, food additives and environmental pollutants. More than 200,000 xenobiotics that exist in our environment are metabolised predominately in the liver. This occurs through the cytochrome P450s enzyme pathways, where the xenobiotics are rendered water soluble, thus being readily eliminated from the body.

We (humans) belong to the ecosystem of this planet. In this sense, we are simply participants in the food chain. While we enjoy diversity from being at the top of the food chain, we experience disadvantages by way of bio-concentration of xenobiotic contaminants (Toxins from the environment). Our exposure to all the environmental toxins including food chemicals, additives, preservatives and drugs that we may take from time to time place a heavier work load on our liver.

The liver is especially important in maintaining a normal blood glucose level. When blood glucose is low, the liver can break down glycogen and release glucose into the bloodstream. The liver also has the ability to convert certain amino acids and lactic acid into glucose, as well as converting other sugars such as fructose and galactose into glucose. When blood glucose is high, as is the case just after eating a meal, the liver converts glucose into glycogen and triglycerides for storage.

The liver is required to carry out Lipid metabolism, as well as Protein metabolism especially the hepatocytes that synthesize most plasma proteins (Such as alpha & beta globulins, albumin, prothrombin & fibrinogen). Our liver is also required for the excretion of bilirubin, which is derived from the heme of the red blood cells. It is absorbed by the liver from the blood and secreted into Bile. Most of the bilirubin in Bile is metabolized in the small intestine by bacteria and eliminated in the feces. In addition, the liver is a prime storage site for certain vitamins, such as Vitamins A, B12, D, E, K and minerals iron & copper. As well as storage of Vitamin D, the liver is involved in synthesizing active forms of Vitamin D. The Kupffer cells of the liver phagocytize aged red and white blood cells and some bacteria.

Our current lifestyles, environmental toxins, consumption of alcohol, food additives and drugs all place our liver under enormous stress and would be impacting on our health and well being.

Some of the symptoms that may indicate that our liver is not detoxifying properly are:

- bowel wind
- constipation
- loose bowels
- dry hair, split ends, dry scalp
- nails splitting or lifting at the edges
- spoon nails
- light coloured stools
- poor appetite
- dermatitis
- psoriasis
- gout
- bad breath
- headaches
- nausea associated with fatty meals
- headaches
- immune weakness
- tiredness & fatigue
- sugar craving
- insomnia
- mood changes
- anxiety
- depression
- skin rashes
- joint pain
- pre-menstrual tension
- high copper levels
- loose bowels
- indigestion
- bloating
- fatty liver
- cirrhosis
- hepatitis
- tiredness
- obesity
- joint pains
- hormonal imbalances
- PMT
- Irritability
- Bile disturbances

So, it is easy to see that with all the toxins we encounter in our environment, plus the toxins that our own bodies produce it is extremely important to maintain a healthy liver.

Cholesterol is synthesized in the liver through the H.M.G CoA reductase pathway. It produces about 700mg of cholesterol per day, while the remainder is derived from the diet. We produce HDL (the 'good' cholesterol), LDL and VLDL cholesterol in our liver, as well as the absorption of dietary cholesterol. All forms of cholesterol are not 'bad' all are necessary, as they are the precursors for steroids in the body (Such as corticosteroid, sex hormones, bile acids and vitamin D.). Problems with cholesterol arise when the ratio between the HDL, LDL & VLDL become unbalanced. Our liver can synthesize cholesterol, but it is also responsible for the removal of cholesterol. Our liver clears around 1g of cholesterol per day, which is eliminated through the bile.

Lecithin is involved with bile production and plays a major role in maintaining solubility of cholesterol. Solubility is important so that we do not develop cholesterol stones in the liver or gall bladder. While our liver can synthesize lecithin, dietary intake is also required.

So, as you can see, it is vitally important to maintain a healthy liver. In doing so, we are also more likely to be in a position of maintaining healthy cholesterol levels.

It is in my opinion, given the level of Xenobiotics in our environment, necessary to supplement with liver nutrients. It would be best to seek a supplement that contains Silybum marianum ext, L. Methionine, L. Glycine, L. Cysteine, L. Glutamine, Inositol, L. Lysine, L. Taurine, Levocarnitine and Glutamic acid, Lecithin, Potassium, Magnesium, Zinc, Selenium, Folic Acid, Pyridoxal-5-Phosphate, B6, B12, Ascorbic Acid, Hesperidin and Rutin. These can all be found in one tablet and would be known as Hepatic Detox.

Given all the work our livers are involved with, it stands to good reason to 'love your liver' and treat it well, so that you may get more out of life.

ACTIVE COMPONENTS

Silybum Marianum (St Mary's Thistle)

St. Mary's Thistle contains three principal biologically active compounds, which are known collectively as Silymarin.

This herb has been found to have a wide range of effects on health. The main target of it's action is the liver, helping to protect this organ against damage from alcohol and other toxins as well helping in cases of Jaundice and cholestasis. However it also has been prescribed for pleurisy, problems with the spleen, dyspepsia, gallbladder problems, as antidote to deathcap mushroom poisoning, depression, psoriasis and hepatitis.

- It protects against lipid peroxidation.
- It improves the antioxidant status in the liver.
- It protects the liver against problems caused by an excessive alcohol intake.
- It helps to protect against liver hepatitis induced by drug and viral hepatitis.
- It protects against alcohol-induced hepatic fibrosis.
- It is beneficial in the prevention or treatment of liver dysfunction in patients undergoing anticancer therapy.
- It has protective actions against some cancers.
- It can reduce plasma VLDL- cholesterol levels.
- It protects against a range of toxins.
- It can modulate the inflammatory immune response.

L Methionine.

Methionine is a sulphur-containing, proteogenic amino acid. It is an essential amino acid. Within the body, the metabolically active form of this amino acid is s-adenosymethionine, which is a methyl group donor in transmethylation reactions. When Methionine is catabolised, cysteine is formed as the liver metabolites.

- Methionine has a protective effect on the liver, as well as protecting the liver against a range of toxic chemicals.
- It is effective in the treatment of alcoholic liver disease; its dietary requirements are increased in liver disease and alcoholic hepatitis.
- It helps to suppress the progression of induced liver fibrosis.
- It may protect liver against alcohol-related carcinogenesis.
- It is effective in the treatment of gallstones.
- It assists the liver with pesticide exposure.
- Methionine is useful in chemical sensitivity.

L Glycine.

Glycine is essential in the diet. As the nitrogen from Glycine can easily be exchanged with other molecules, Glycine can be used as a starting point for the synthesis of other amino acids.

- It assists in the detoxification through its promotion of glutathione availability.
- It has a protective effect on the liver in a wide range of toxins and chemical sensitivity.
- It has a role in protecting tissues against ischemia
- It has a protective effect in the liver inflammatory diseases.
- It protects against shock caused either by blood loss or endotoxin, reduces alcohol levels in the stomach and recovery from alcoholic hepatitis.
- It helps to protect the liver damage caused by hepatotoxic drugs.
- It can normalize fatty acid levels in the tissues.
- It reduces the free radical-mediated oxidative stress in the erythrocyte membrane, plasma and hepatocytes in alcohol-induced liver injury.
- It helps protect against obstructive jaundice.

L Cysteine.

Cysteine, an amino acid is a key molecule in sulphur metabolism in the body. Cysteine also has a sparing effect on Methionine for protecting mucosa linings and helping to protect the body against free radicals.

- It helps to protect the body against free radicals.
- It is involved in the production of glutathione within the liver, which reduce the oxidative damage in hepatocytes.
- It protects against toxic substances in the liver, including those from cigarette smoke.
- It protects against liver cholestasis.
- It reduces the oxidative stress of hepatitis C in the liver.
- It helps the liver in chemical sensitivity.
- It is useful in helping to protect the liver against liver disease.
- It is involved with reducing heavy metal toxicity.

L Glutamine.

Glutamine is one of the proteogenic amino acids. It can be synthesized from glutamic acid, a reaction that is important in the detoxification reaction that helps to remove the toxic ammonia from the body. Glutamine is an intermediate nitrogen, allowing its nitrogen to be transferred to other molecules in transamination reactions.

- It removes ammonia from the hepatocytes (liver).
- It is important for maintenance of cell function.
- It promotes storage of glycogen in the muscle.
- It is necessary for normal intestinal morphology and function.
- Decreased glutamine levels are associated with catabolic stress and are associated with increased susceptibility to infection.
- It protects the liver against toxic substances.
- It is useful in the treatment of chronic hepatitis.
- Anyone with defective liver function could benefit from supplementation with glutamine
- It is one of the building blocks for glutathione, important in the body's antioxidant defence.
- It helps in chemically sensitive individuals.


L Lysine

Lysine is an essential amino acid, one of the groups known as proteogenic. The biological active from this molecule is L-lysine, Advanced Medicine Australia Hepatic Detox only uses L amino acid.

- A deficiency of this amino acid can lead to a number of stresses in the tissues.
- L-lysine can suppress recurrent herpes simplex infection.
- It enhances calcium absorption from the intestines and reduces the losses of this mineral through the kidney.
- It helps the prevention of cardiovascular disease and osteoporosis.
- It has a protective action on the liver following haemorrhage.
- It helps in detoxification of lead.



get more
out of life!



L Taurine.

Taurine is an amino acid. It is a sulphur-containing molecule found in bile in combination with cholic acid. In the bile Taurine is one of the agents that help to emulsify lipids (fats), assisting their absorption and transport. Taurine assists in regulating the level of bile salt in the gall bladder, reducing the incidence of gall stones.

Taurine is useful in the treatment of fatty liver. In a study of fatty liver in children with simple obesity, it was found that taurine supplements in the diet were effective in treating their fatty liver. The authors suggested that taurine administration may be helpful as an adjuvant therapy for fatty liver.

Taurine helps to protect against liver fibrosis which is the result of excessive protein deposition by hepatic stellate cells. In a study it was found that taurine supplementation prevented the development of fibrosis of the liver.

Taurine seems to protect the liver against some deleterious changes during hepatocarcinogenesis. This was discovered in a study of rats fed taurine supplements, and where hepatocarcinogenesis was induced. In this study, taurine not only inhibited lipid peroxidation in the liver, but also altered defence enzymes and protected the liver against membrane disintegration during the hepatocarcinogenesis (You JS, Chang KJ, 1998).

Taurine is one of several amino acids that have been found to be essential in disease states, and to form substrate in the stressed patient for anabolic processes in the liver, immune system, and at injured sites (Soeters PB et al, 2004).

- It protects the tissues and liver tissue against oxidative stress.
- Its levels are reduced when the liver is diseased or stressed.
- It is useful in the treatment of fatty liver.
- It protects the liver against liver fibrosis.
- It is involved with the detoxification of chemical toxins.
- Involved in bile synthesis.
- It facilitates the passage of calcium, potassium, sodium, and magnesium ions in and out of cells.
- It is useful in gall bladder disease.
- It protects against cholesterol gallstone formation.
- It protects the liver against some deleterious changes during hepatocarcinogenesis.
- Taurine helps to protect the liver against fatty liver.

L Carnitine.

Carnitine is important for all tissues that have high-energy demand like the liver. In a study of patients with cirrhosis of the liver, Lapinski T, Grzeszczuk A (2003) found that carnitine lowered serum ammonia concentration and lipid metabolism in the liver. This reinforces the role of carnitine as a protective agent for the liver. Carnitine facilitates the removal from mitochondria of short-chain and medium-chain fatty acids that accumulate as a result of normal and abnormal metabolism. One other function of Carnitine is to remove potentially toxic acyl groups from cells by promoting the formation of acylcarnitine from acyl-CoA. This system is important for removing the toxic acyl groups.

- It removes potentially toxic acyl groups from cells.
- Carnitine has been used in senile dementia, metabolic nerve diseases, HIV infection, tuberculosis, myopathies, cardiomyopathies, renal failure and anemia.
- It lowers serum ammonia concentrations and improves lipid metabolism in the liver.
- It protects against hepatic encephalopathy.
- It prevents lipid preoxidation.
- It reduces lipid accumulation in the liver.
- A deficiency in carnitine has been implicated in the development of non-alcoholic related fatty liver disease.



hepatic detox

L Glutamic acid

Glutamic acid has its main role is in the transport of potassium in the brain, and in the detoxification of ammonia in the brain.

- In the liver, it modulates the rate of ammonia detoxification into urea.
- It promotes glycogen storage.
- It is needed for the synthesis of glutathione within the liver, a critical antioxidant for the body.
- It protects against free radicals in the liver.
- It is needed in alcoholics and cirrhotics and in acute liver failure.
- Its ability to reduce ammonia levels in the brain protects against hepatic encephalopathy.

Inositol

Inositol is necessary for normal cellular proliferation (cell division). Inositol deficiency may promote fatty liver disease. The impaired release of plasma lipoprotein increases fatty acid mobilization from adipose tissue and enhanced fatty acid synthesis in the liver. Inositol deficiency has causative factors in the occurrence of hepatic triacylglycerol accumulation. Very little is known about calcium signaling through the Inositol receptor, which is a mechanism in cells to regulate their function. However, it is known that this receptor is lost from cholangiocytes in cholestatic liver disease and this may have significance in cholestasis. In turn, this suggests that any deficiency in Inositol may have a negative impact on the development of cholestasis.

- A potent ability to reduce hepatic lipid levels and regulate blood lipids.
- It protects the liver against toxins.
- It protects the liver against fatty liver.
- It is necessary for normal cellular proliferation (cell division).
- It has a protective effect against cancers in the liver and lungs and possibly also in the mammary gland and colon.
- It may help in the treatment for diabetic neuropathy.

Lecithin

Lecithin is a phospholipid, specifically a glycerophospholipid, which is derived from phosphoric acid. Lecithin supplementation has a protective effect on the liver against fatty liver. In a clinical trial of patients with fatty liver, where the fat deposits in the hepatocytes were disruptive of their function, it was found that lecithin supplementation minimized this insult on the hepatocytes (Turecky L et al, 2003). Lecithin helps to prevent gallstone formation by stimulating bile formation and biliary lipid secretion, particularly of cholesterol. Lecithin has its effect of solubilisation of cholesterol within the bile, preventing the formation of gallstones.

- It has significant properties that help to protect the tissues against oxidative damage.
- It protects against fatty liver disease.
- It decreases oxidative stress in the liver by helping to restore glutathione levels.
- It helps prevent and reverse hepatic (liver) fibrosis.
- It helps to prevent gallstone formation.
- It has beneficial effect on the vascular system through modification of hepatic cholesterol homeostasis and lipoprotein metabolism.
- Lecithin which is a good source of phosphatidylcholine.

Potassium

The regulation of potassium levels in the body is of crucial importance. One of the ways potassium is regulated is through an insulin-mediated uptake and release from the liver and muscles, helping to regulate the distribution of this molecule between the intercellular and extracellular fluid compartments (Decaux G et al, 1988). In studies on humans, these authors found that liver damage decreased uptake of potassium into the liver, despite there being more than adequate amounts of insulin available to promote this. Fasting also decreases the levels of potassium in the liver. It has been found in studies that potassium may help to protect the liver from tumor necrosis factor.

- Fasting decreases the levels of potassium in the liver.
- It helps to protect the liver from tumor necrosis factor.
- It is one of the dietary measures that have been suggested as a way of reducing the risk of Alzheimer disease.

Magnesium.

Magnesium, a metal, is an essential micronutrient for humans. Magnesium deficiency has been noted when the liver is under environmental stress. It has been proposed that this could result from an inadequate dietary intake of magnesium, poor absorption of magnesium from the gut, or from loss of magnesium as a result of the use of natriuretic drugs (Oralewska B et al, 1996). These authors found deficiencies in 30% of children that they studied. It was further suggested that the fall in magnesium levels in the blood is a reflection of the extent of the damage to the liver cells. In another study with magnesium the authors found that insulin sensitivity was observed and insulin secretion was reduced with magnesium deficiencies. It also was observed that when magnesium supplementation was given, the insulin resistance restored to normal in the liver. They proposed that it was a result of magnesium having an early action on the interaction of insulin with the hepatic cells. These authors, (Rozycky J, Lewicki Z, 1978), found that low magnesium levels may promote the deposition of fat in the liver. Magnesium deficiency results in both proliferation and enlargement of Ito cells in the liver. These authors proposed that the lipid accumulation resulted from reduced levels of lipid metabolizing and transporting enzymes, which in turn was a consequence of the lack of magnesium needed in the enzyme structure.

- Liver stress may result in magnesium deficiency.
- Low magnesium levels may promote the deposition of fat in the liver.
- Magnesium supplementation is needed with insulin resistance within the liver.
- Magnesium deficiency may cause liver disease.

Aspartate

Aspartate in hepatic detox is found as magnesium and potassium aspartate. Aspartate is an amino acid. The main role in the liver is in the urea cycle and in purine and pyrimidine biosynthesis. Aspartate is important in assisting the liver to cope with toxins. In particular aspartate helps the liver clear toxic ammonia by promoting the formation of urea.

- It assists the liver to cope with toxins.
- It improves urea production, helping to detoxify the liver of ammonia.
- It may have clinical usefulness in patients with acute or chronic liver failure.

Zinc

Zinc is a metal that is essential for growth and development, as it is an important component of many enzymes. Zinc deficiencies increase the susceptibility to some viral infections that attack the liver. In one study it was found that nutritional deficiencies, such as vitamin E, Folic Acid, Vitamin B6, Zinc, and selenium resulted in alteration of the immune system that eventually increase susceptibility to certain virus infection such as hepatitis B virus and hepatitis C virus (Poschl G, Seitz HK, 2004). In another study it linked zinc deficiency to liver cirrhosis and diabetes mellitus. Zinc supplementation increased glucose disposal. It was proposed by these authors that zinc deficiency is linked between liver cirrhosis and "liver" diabetes mellitus (Grungreiff K, Reinhold D, 2005).

Zinc helps to protect against toxins like lead and cadmium, and Zinc also has a balancing effect with copper when in excess. Zinc and magnesium have a synergistic effect within the liver for many enzyme functions.

- Its deficiency increases the susceptibility to some viral infections that attack the liver.
- Zinc deficiency is a characteristic of both liver cirrhosis and diabetes mellitus.
- Diets deficient in zinc have an adverse impact on the liver.
- Zinc deficiency depresses glycogen catabolism in the liver.
- It helps to protect against some toxins.
- Alcohol consumption can lead to liver damage and adversely affect zinc levels in the liver.
- It can raise glutathione peroxidase levels in the liver, reducing the risk of lipid peroxidation in the hepatocytes and helping to maintain the integrity of the cell membranes of the hepatocytes.
- It protects against some cancers.

Selenium

Selenium is an essential nutrient for humans. It is an essential component of the enzyme glutathione peroxidase, which helps protect against oxidative damage in the liver and particularly in the membranes of the red corpuscles.

Selenium is essential for normal liver function. One role in the liver is to combine with the amino acid cysteine to form selenocysteine, which is then incorporated into a class of proteins known as selenoproteins, which have both antioxidant and selenium transport functions. It has been demonstrated that inadequate levels of selenoproteins in the liver can result in hepatocellular degeneration, liver necrosis and death. Other papers report a protective effect of selenoprotein against toxins that can cause damage to the liver. The biochemical and physicochemical properties of selenium in the unique redox characteristics of selenocysteine and its use in antioxidant enzymes. In this context of a redox reaction is reduction of reactive oxygen metabolites by glutathione peroxidases, helping to maintain membrane integrity, reducing the oxidative damage to lipids, lipoprotein and DNA. Selenium has structural and enzymatic roles, and also influences a number of endocrine processes, most notably those involved in thyroid hormones synthesis and metabolism.

Selenium is needed for the proper functioning of the immune system, a role in viral suppression, AIDS, and is implicated in delaying the aging process. Its deficiency has been linked to a number of disorders such as heart disease, diabetes and diseases of the liver. It is required for sperm motility and may reduce the risk of miscarriage. Selenium supplementation has moved from the realm of correcting nutritional deficiencies to one of pharmacological intervention, especially in the clinical domain of cancer chemoprevention.

Selenium protects against aluminium toxicity; something that is useful as this metal can be neurotoxic. It also protects against cadmium poisoning in the liver as well as protects against lead toxicity.

Selenium also has a protective role in the liver against a wide range of heavy metals. It was found that selenium within the liver helped to protect against mercury, silver, copper and cadmium. It had its effect in several locations within the cells. Mercury and silver preferentially accumulates in the nuclear, lysosomal and mitochondrial fraction of the hepatocytes, whereas copper and cadmium accumulates mainly in the cytosol. Cadmium impairs the liver of detoxification.

- It can positively influence the health status, symptomatic presentation and life span whose detoxification systems are compromised.
- It is essential for liver function.
- It protects against disease and aging.
- Exercise increases the requirement for selenium.
- It promotes antioxidant activity in the tissues.
- It protects against a wide range of toxins, including heavy metals.
- It protects against actions of the prednisolone and some chemotherapeutic agents in the liver.
- It has a protective action in alcoholic liver disease.
- It has a protective action against some cancers.
- It protects against some viral infections that attack the liver.



love
your
liver

Folic acid

Also known as folate, folic acid is essential in the body. At least one-half of the folate stores in the body are found in the liver, with the level of these stores being influenced by the dietary intake of this vitamin. Folate deficiency does remove some protection from DNA strands and this can promote cancer within the liver. This is because folate, together with other molecules such as Methionine and vitamin B12, play an important role in DNA methylation and this methylation influences the expression of genes. The enzyme responsible for this methylation is methylenetetrahydrofolate reductase, which is responsible for the de novo synthesis of methyl groups. Its stated that diminished activity of this enzyme, which is derived from folic acid, is associated with an increased risk of developing atherosclerosis, neurological disorders, birth defects and may lead to liver cancer.

Folic acid, together with vitamin B12, B6 and the activated form of B6 pyridoxine-5- phosphate has the role of reducing the effects of homocysteine in the blood plasma. The supplementation of these vitamins can reduce levels of low-density lipoprotein in cases of high hyperhomocysteinaemia, thereby reducing risk of atherosclerosis and possibly also diabetes and alcoholic liver disease.

Folic acid helps to protect the liver against some toxins. By providing folate it was found that it had important influence on toxins. It has effects on the glutathione level within the liver, an important antioxidant for the liver defense.

- Deficiencies in folic acid intake increase the susceptibility to some viral infections that attack the liver.
- Folic acid deficiency promotes the carcinogenic consequences of alcohol metabolism and cigarette smoking.
- It improves normal liver morphology.
- It helps the treatment or prevention of atherosclerosis and other condition in the people with certain genetic disorders.
- It helps to protect the liver against some toxins.
- It is a potent scavenger of free radicals and can significantly inhibit microsomal lipid peroxidation.
- It may have a role in the prevention of Alzheimer disease.
- It may help in the prevention of birth defect.

Vitamin B6 and Pyridoxal-5-Phosphate

Vitamin B6 is also known as Pyridoxine. Vitamin B6 and Pyridoxal-5-Phosphate is a water-soluble vitamin. The concentration is mainly found in the liver and kidneys. In the body this vitamin is important as a coenzyme in amino acid metabolism. Additionally pyridoxine and pyridoxal-5-phosphate is involved in the release of glucose from glycogen in the liver.

In patients with fatty liver (cirrhosis) there is a decreased metabolism in the liver of vitamin B6 to pyridoxal-5-phosphate (the active coenzyme form of B6), which is reflected by low levels of this molecule in the plasma of these patients. However they found that oral supplementation with pyridoxine and pyridoxal-5-phosphate restored levels to normal range, indicating a supportive effect of the vitamin in the liver. In a related study, plasma levels of pyridoxal-5-phosphate were found to be significantly lower than normal in patients with decompensated cirrhosis or subacute hepatic necrosis, with no significant difference in plasma pyridoxal-5-phosphate levels between those with liver disease. All patients responded to supplements of pyridoxal-5-phosphate and vitamin B6. Vitamin B6 is involved with reducing the effects of homocysteine.

- Through its methylating ability it may prevent the development of some cancers.
- It mobilizes liver glycogen, resulting in an increase in serum glucose levels.
- It has an impact on the metabolism of both lipids and proteins.
- Deficiency states can be caused by alcohol and fasting.
- Deficiencies of B6 may have a role in causing fatty liver.
- Deficiencies in vitamin B6 intake increase the susceptibility to some viral infection that attack the liver.
- Vitamin B6 reduces the effect of homocysteine.

Vitamin B12

Also known as cobalamin, or extrinsic factor, vitamin B12 is one of the water-soluble B group vitamins. This vitamin was the first naturally occurring biologically active molecule found to require cobalt as a part of its molecular structure. In nature bacteria synthesize this vitamin, while green plants contain none of the vitamin. The main dietary sources are egg yolk and milk, which means that some strict vegetarians may become deficient in this vitamin. Absorption of this vitamin from the gut requires the presence of a molecule referred to as intrinsic factor, which resides in the gastric mucosa. This factor forms a pepsin resistant combination with the vitamin B12, allowing it to be absorbed in the lower intestine. Pernicious anemia develops when intrinsic factor is defective. Vitamin B12 binds to other groups to have a range of biological effects in the body. Cyanocobalamin (vitamin B12) is needed to prevent the development of pernicious anemia. Vitamin B12 forms a coenzyme needed for the methylation of RNA, a process that is important in protein synthesis. Vitamin B12 and folate are needed as coenzymes for DNA methylation, which are methyl donors during DNA synthesis. DNA methylation is an intracellular process that can influence the expression of some genes and can therefore effect cellular metabolism. The transmethylation metabolic pathways, which provide or removed the methyl groups closely, interconnect the important methyl group molecules; chorine, Methionine, methyltetrahydrofolate, vitamin B6 and vitamin B12.

Abnormal DNA methylation has been associated with liver cancers (Brunaud Let al, 2003). This means that deficiencies in any of the major dietary sources of methyl groups (such as Methionine and chorine), or a deficiency in vitamin B12 which is needed as a coenzyme in the methylation process, may result in the development of liver cancers as well as other cancers. Deficiencies in vitamin B12 disrupt metabolism within the liver. In a study of livers it was found that this disrupted normal glucose and glutamic acid metabolism, possibly though an inhibition on the citric acid cycle (Toyoshima S et al, 1996). A deficiency in B12 results in decreased folate incorporation in tissues of the liver. It also results in depressed synthesis in Methionine and lecithin. Lecithin is an important protective molecule in the liver. These changes can result in alterations in membrane lipid composition and therefore in cellular function. Deficiencies in a range of nutrients including vitamin B12 can result in hyperhomocysteinemia, which is a component of several disorders including cardiovascular disease, neurodegeneration, diabetes and liver disease. Hyperhomocysteinemia increases production of intracellular superoxide anion causing oxidative stress and induces endoplasmic reticulum stress, which can result in apoptosis, fat accumulation and inflammation.

- Through its methylating ability it may prevent the development of some cancers.
- Liver disease can result in a loss of vitamin B12 from the liver.
- Deficiencies in vitamin B12 disrupt metabolism within the liver.
- Some toxins and disease states result in disturbances in vitamin B12 metabolism.
- It protects against some of the action of toxins on the liver.
- Its supplementation may have a role in the prevention of Alzheimer disease.
- B12 supplementation may help reduce hyperhomocysteinemia.

Hepatic Detox is a comprehensive liver support nutrient required for good liver function and helps support detoxification.

Ascorbic acid

Also known as vitamin C, ascorbic acid is a water-soluble vitamin essential for health. As it cannot be synthesized in the body, it must be ingested. This vitamin is important in the body in many hydroxylation reaction (such as that involving the conversion of the amino acid proline into collagen). It is critical for maintaining the structure of the walls of blood vessels because of its role in collagen biosynthesis. Because vitamin C is a potent antioxidant, it helps to protect all tissues in the body. This is because many toxic agents exert their effects in the body by increasing the levels of damaging free radicals in the tissues and antioxidants are able to scavenge these free radicals to protect the tissues. Vitamin C assists in the mitigation of heavy metal toxicity. In one study it looked at cadmium and arsenic, two particularly toxic chemicals, it was found that ascorbic acid protects against the effects of arsenic on mitochondria. Vitamin C also appears to be protective against lead toxicity. The liver is a target organ for 3,4-methylenedioxymethamphetamine (MDMA, 'Ecstasy'), resulting in toxicity in that organ when this drug is taken. This toxicity appears to result from impaired antioxidant defense systems, but it can be alleviated somewhat by increased levels of antioxidants such as vitamin C. Evidence supporting vitamin C on the liver was provided in a good study on ischaemic livers, where the vitamin C scavenging of free radicals helped to reduce injury to liver tissue. Similar results have been found in vitamin C and its effects to protect the liver following damage by toxicants. One of the consequences of oxidative stress is the induction of collagen synthesis, which in the liver could lead to reduced function of the hepatocytes. This fibrosis can be induced with alcohol. The authors concluded that a mixture of antioxidants, including vitamin C, is needed.

- It protects against heavy metal toxicity.
- It protects the liver against oxidative stress.
- It reduces the chance of fibrosis developing in the liver.
- It protects against the effects of alcohol on the liver.
- It protects against atherosclerosis and other diseases by preventing abnormal lipid metabolism.

Hesperidin

Hesperidin is a flavone. These are plant pigments that contain flavone ring system. Hesperidin is known as a bioflavonoid, Hesperidin has a range of beneficial effect in the body. Because of their molecular structure, the bioflavonoids are antioxidants, and therefore help to protect tissues against oxidative damage. Indeed the pharmacological antioxidant properties of Hesperidin are similar to those of vitamin E. Dietary supplementation with Hesperidin has demonstrated the lowering of hepatic triglyceride content, as well as plasma cholesterol levels. The ability of Hesperidin to lower triglyceride levels in the liver has been demonstrated in studies. It was shown that dietary Hesperidin reduced the hepatic triglyceride accumulation induced by orotic acid (in rats) and this was associated with the reduced activity of the hepatic microsomal triglyceride synthetic enzyme, phosphatidate phosphohydrolase. In studies on diabetes, Hesperidin has been found to reduce oxidative stress in the tissues, demonstrating further the antioxidant properties of this bioflavonoid.

In nature, bioflavonoids are found together with vitamin C. Vitamin C and the bioflavonoids have a bewildering array of application which makes them appear like the old medicine man's magic, something which does not endear them to the orthodox thinkers. Dr Robbins of the university of Florida, investigated the antithrombic activity of bioflavonoids for about 6-7 years. He had research showing that bioflavonoids have anti-adhesive effects on blood cells. That is, they prevent blood cells from clumping together and forming blood cots, which can cause massive damage to the circulation through the heart, lungs, brain and liver. His experiments showed that bioflavonoids were indeed potent antithrombins.

- It lowers hepatic triglyceride content, as well as plasma cholesterol levels.
- It reduces oxidative stress in the tissues.
- It offers some protection against the early stages of diabetes mellitus and its complications.
- It is found in nature together with vitamin C and has a synergistic effect together with vitamin C.

Rutin

Rutin, like Hesperidin, is a flavone, also known as a bioflavonoid. Rutin is the most potent scavenger of superoxide anions, more potent than its near relative, Hesperidin. Rutin, together with other antioxidants, is important in protecting us from disease and aging. In situations where there is inadequate vitamin E (which normally is a potent antioxidant), Rutin and other flavonoids can take over from vitamin E as a major antioxidant in the liver. Rutin has a substantial protective effect on the liver. Dietary supplementation with Rutin has demonstrated to increase the antioxidant capacity of the liver. This study has been supported in a study of human volunteers who took Rutin supplements over six weeks. These studies indicate that Rutin could be a useful supplement for anyone whose liver is under assault by toxic agents. When there is an iron overload in the body, Rutin has been found to be protective possibly through the formation of inactive iron-Rutin complexes. In rat models that were studied, Rutin was found to substantially reduce free radical production in lipid microsomes.

- It protects against disease and aging.
- It is a potent antioxidant.
- It protects against oxidative damage to DNA and proteins.
- It has a substantial protective effect on the liver.
- It protects the liver against a range of toxic chemicals.
- It helps to protect against some carcinogens, including those in cigarette smoke.
- It protects against an iron overload in the body.

Suggested Uses for Hepatic Detox:

- EXPOSURE TO HEAVY METALS & ENVIRONMENTAL TOXINS
- HIGH COPPER LEVELS
- CONSTIPATION
- LOOSE BOWELS
- INDIGESTION
- BLOATING
- POOR APPETITE
- FOOD ALLERGIES
- BAD BREATH
- PSORIASIS
- NAUSEA ASSOCIATED WITH FATTY MEALS
- FATTY LIVER
- CIRRHOSIS
- HEPATITIS
- HANGOVERS
- ALCOHOLICS
- BOWEL WIND
- HEADACHES / MIGRANES
- SUGAR CRAVINGS
- SKIN PROBLEMS
- DERMATITIS
- ACNE
- SKIN RASHES
- TIREDNESS
- FATIGUE
- INSOMNIA
- IMMUNE WEAKNESS
- DIET HIGH IN PROCESSED FOODS
- AGING PROCESS
- OBESITY
- WEIGHT LOSS
- STRESS
- LIGHT COLOURED STOOLS
- HYPERCHOLESTEROLEMIA
- DRY HAIR
- HAIR - SPLIT ENDS
- DRY SCALP
- GOUT
- JOINT PAINS
- SPOON NAILS
- NAILS SPLITTING OR LIFTING AT THE EDGES
- HORMONAL IMBALANCES
- PMT
- USERS OF ORAL CONTRACEPTIVE
- IRRITABILITY
- ANGER
- DEPRESSION
- ANXIETY
- DRUG USE e.g. PARACETAMOL
- BILE DISTURBANCES

References

References for Silybum Marianum

Ahmed B, Khan SA, Alam T (2003) Synthesis and antihepatotoxic activity of some heterocyclic compounds containing the 1,4-dioxane ring system. *Pharmazie*. 2003 Mar;58(3):173-6

Bean P (2002) The use of alternative medicine in the treatment of hepatitis C. *Am Clin Lab*. 2002 May;21(4):19-21

Beckmann-Knopp S et al (2000) Inhibitory effects of silibinin on cytochrome P-450 enzymes in human liver microsomes. *Pharmacol Toxicol*. 2000 Jun;86(6):250-6

Gebhardt R (2003) Antioxidative, antiproliferative and biochemical effects in HepG2 cells of a homeopathic remedy and its constituent plant tinctures tested separately or in combination. *Arzneimittelforschung*. 2003;53(12):823-30

Giese LA (2001) Milk thistle and the treatment of hepatitis. *Gastroenterol Nurs*. 2001 Mar-Apr;24(2):95-7

Hernandez V (1995) Liver function and HIV-1 infection. *STEP Perspect*. 1995 Summer;7(2):13-5

Jacobs BP et al (2002) Milk thistle for the treatment of liver disease: a systematic review and meta-analysis. *Am J Med*. 2002 Oct 15;113(6):506-15

Johnson VJ et al (2003) Physiological responses of a natural antioxidant flavonoid mixture, silymarin, in BALB/c mice: III. Silymarin inhibits T-lymphocyte function at low doses but stimulates inflammatory processes at high doses. *Planta Med*. 2003 Jan;69(1):44-9

Kohno H et al (2002) Silymarin, a naturally occurring polyphenolic antioxidant flavonoid, inhibits azoxymethane-induced colon carcinogenesis in male F344 rats. *Int J Cancer*. 2002 Oct 10;101(5):461-8

Kravchenko LV, Morozov SV, Tutelyan VA (2003) Effects of flavonoids on the resistance of microsomes to lipid peroxidation in vitro and ex vivo. *Bull Exp Biol Med*. 2003 Dec;136(6):572-5

Ladas EJ, Kelly KM (2003) Milk thistle: is there a role for its use as an adjunct therapy in patients with cancer? *J Altern Complement Med*. 2003 Jun;9(3):411-6

Laekeman G, De Coster S, De Meyer K (2003) St. Mary's Thistle: an overview. *J Pharm Belg*. 2003;58(1):28-31

Lieber CS et al (2003c) Silymarin retards the progression of alcohol-induced hepatic fibrosis in baboons. *J Clin Gastroenterol*. 2003 Oct;37(4):336-9

Rambaldi A et al (2005) Milk thistle for alcoholic and/or hepatitis B or C virus liver diseases. *Cochrane Database Syst Rev*. 2005 Apr 18;(2):CD003620

Saller R, Meier R, Brignoli R (2001) The use of silymarin in the treatment of liver diseases. *Drugs*. 2001;61(14):2035-63

Schumann J et al (2003) Silibinin protects mice from T cell-dependent liver injury. *J Hepatol*. 2003 Sep;39(3):333-40

Siddiqui U, Weinschel EH, Bini EJ (2004) Prevalence and predictors of herbal medication use in veterans with chronic hepatitis C. *J Clin Gastroenterol*. 2004 Aug;38(7):605-10

Skottova N et al (2004) Phenolics-rich extracts from Silybum marianum and Prunella vulgaris reduce a high-sucrose diet induced oxidative stress in hereditary hypertriglyceridemic rats. *Pharmacol Res*. 2004 Aug;50(2):123-30

Svagera Z et al (2003) Plasma lipoproteins in transport of silibinin, an antioxidant flavonolignan from Silybum marianum. *Phytother Res*. 2003 May;17(5):524-30

Tanamly MD et al (2004) Randomised double-blinded trial evaluating silymarin for chronic hepatitis C in an Egyptian village: study description and 12-month results. *Dig Liver Dis*. 2004 Nov;36(11):752-9

Yanai Y et al (2002) Dietary silymarin suppresses 4-nitroquinoline 1-oxide-induced tongue carcinogenesis in male F344 rats. *Carcinogenesis*. 2002 May;23(5):787-94

Zuber R et al (2002) Effect of silybin and its congeners on human liver microsomal cytochrome P450 activities. *Phytother Res*. 2002 Nov;16(7):632-8

References for Methionine

Brunaud L et al (2003) Effects of vitamin B12 and folate deficiencies on DNA methylation and carcinogenesis in rat liver. *Clin Chem Lab Med*. 2003 Aug;41(8):1012-9

Glazenburg EJ et al (1983) Effects of variation in the dietary supply of cysteine and methionine on liver concentration of glutathione and "active sulfate" (PAPS) and serum levels of sulfate, cysteine, methionine and taurine: relation to the metabolism of acetaminophen. *J Nutr*. 1983 Jul;113(7):1363-73

Halsted CH et al (2002b) Folate deficiency disturbs hepatic methionine metabolism and promotes liver injury in the ethanol-fed micropig. *Proc Natl Acad Sci U S A*. 2002 Jul 23;99(15):10072-7. Epub 2002 Jul 16

Halsted CH, Villanueva JA, Devlin AM (2002) Folate deficiency, methionine metabolism, and alcoholic liver disease. *Alcohol*. 2002 Jul;27(3):169-72

Lee TD et al (2004) Abnormal hepatic methionine and glutathione metabolism in patients with alcoholic hepatitis. *Alcohol Clin Exp Res*. 2004 Jan;28(1):173-81

Lieber CS (1993) Biochemical factors in alcoholic liver disease. *Semin Liver Dis*. 1993 May;13(2):136-53

Lieber CS (2000) Alcoholic liver disease: new insights in pathogenesis lead to new treatments. *J Hepatol*. 2000;32(1 Suppl):113-28

Lieber CS (2002) S-adenosyl-L-methionine: its role in the treatment of liver disorders. *Am J Clin Nutr*. 2002 Nov;76(5):1183S-7S

Loguercio C et al (1999) Ethanol consumption, amino acid and glutathione blood levels in patients with and without chronic liver disease. *Alcohol Clin Exp Res*. 1999 Nov;23(11):1780-4

Matsui H et al (2004) Sulfur-containing amino acids attenuate the development of liver fibrosis in rats through down-regulation of stellate cell activation. *J Hepatol*. 2004 Jun;40(6):917-25.

Niculescu MD, Zeisel SH (2002) Diet, methyl donors and DNA methylation: interactions between dietary folate, methionine and choline. *J Nutr*. 2002 Aug;132(8 Suppl):2333S-2335S

Nocianitri KA et al (2002) Influence of dietary methionine level on the liver metallothionein mRNA level in rats. *Biosci Biotechnol Biochem*. 2002 Nov;66(11):2465-70

Patrick L (2003) Toxic metals and antioxidants: Part II. The role of antioxidants in arsenic and cadmium toxicity. *Altern Med Rev*. 2003 May;8(2):106-28

Poirier LA (2002) The effects of diet, genetics and chemicals on toxicity and aberrant DNA methylation: an introduction. *J Nutr*. 2002 Aug;132(8 Suppl):2336S-2339S

Poschl G et al (2004) Alcohol and cancer: genetic and nutritional aspects. *Proc Nutr Soc*. 2004 Feb;63(1):65-71

Reicks MM, Hathcock JN (1984) Effects of dietary methionine and ethanol on acetaminophen hepatotoxicity in mice. *Drug Nutr Interact*. 1984;3(1):43-51

Ronchi VP et al (2004) The mouse liver content of carbonic anhydrase III and glutathione S-transferases A3 and P1 depend on dietary supply of methionine and cysteine. *Int J Biochem Cell Biol*. 2004 Oct;36(10):1993-2004

Villanueva JA, Halsted CH (2004) Hepatic transmethylation reactions in micropigs with alcoholic liver disease. *Hepatology*. 2004 May;39(5):1303-10

References for Glycine

Bruck R (2003) Glycine modulates cytokine secretion, inhibits hepatic damage and improves survival in a model of endotoxemia in mice. *Liver Int*. 2003 Aug;23(4):276-82

Fang HQ et al (2003) Effects of glycine on plasma and liver tissue changes of TNF-alpha, ET-1 and nitric oxide contents in rats with obstructive jaundice. *World J Gastroenterol*. 2003 Oct;9(10):2374-6

Hall JC (1998) Glycine. *JPEN J Parenter Enteral Nutr*. 1998 Nov-Dec;22(6):393-8

Kretzschmar M (1996) Regulation of hepatic glutathione metabolism and its role in hepatotoxicity. *Exp Toxicol Pathol*. 1996 Jul;48(5):439-46

Lauterburg BH (2002) Analgesics and glutathione. *Am J Ther*. 2002 May-Jun;9(3):225-33

Loguercio C et al (1999) Ethanol consumption, amino acid and glutathione blood levels in patients with and without chronic liver disease. *Alcohol Clin Exp Res*. 1999 Nov;23(11):1780-4

Matilla B et al (2002) Glycine: a cell-protecting anti-oxidant nutrient. *Nutr Hosp*. 2002 Jan-Feb;17(1):2-9

Qu W et al (2002) Glycine blocks the increase in intracellular free Ca²⁺ due to vasoactive mediators in hepatic parenchymal cells. *Am J Physiol Gastrointest Liver Physiol*. 2002 Dec;283(6):G1249-56. Epub 2002 Aug 14

Senthilkumar R, Nalini N (2004) Effect of glycine on tissue fatty acid composition in an experimental model of alcohol-induced hepatotoxicity. *Clin Exp Pharmacol Physiol*. 2004 Jul;31(7):456-61

Senthilkumar R, Nalini N (2004b) The potential beneficial effect of glycine on the carbohydrate moieties of glycoproteins in an experimental model of alcohol-induced hepatotoxicity. *Med Food*. 2004 Spring;7(1):108-13

Senthilkumar R, Nalini N (2004c) Glycine prevents hepatic fibrosis by preventing the accumulation of collagen in rats with alcoholic liver injury. *Pol J Pharmacol*. 2004 Jan-Feb;56(1):121-8

Senthilkumar R, Nalini N (2004b) The potential beneficial effect of glycine on the carbohydrate moieties of glycoproteins in an experimental model of alcohol-induced hepatotoxicity. *Med Food*. 2004 Spring;7(1):108-13

Senthilkumar R, Nalini N (2004c) Glycine prevents hepatic fibrosis by preventing the accumulation of collagen in rats with alcoholic liver injury. *Pol J Pharmacol*. 2004 Jan-Feb;56(1):121-8

Senthilkumar R, Sengottavelan M, Nalini N (2004) Protective effect of glycine supplementation on the levels of lipid peroxidation and antioxidant enzymes in the erythrocyte of rats with alcohol-induced liver injury. *Cell Biochem Funct*. 2004 Mar-Apr;22(2):123-8

Senthilkumar R, Viswanathan P, Nalini N (2003) Glycine modulates hepatic lipid accumulation in alcohol-induced liver injury. *Pol J Pharmacol*. 2003 Jul-Aug;55(4):603-11

Senthilkumar R, Viswanathan P, Nalini N (2004) Effect of glycine on oxidative stress in rats with alcohol induced liver injury. *Pharmazie*. 2004 Jan;59(1):55-60

Thurman RG et al (1998) Kupffer cell-dependent reperfusion injury in liver transplantation: new clinically relevant use of glycine. *Langenbecks Arch Chir Suppl Kongressbd*. 1998;115:185-90

Wheeler MD et al (1999) Glycine: a new anti-inflammatory immunonutrient. *Cell Mol Life Sci*. 1999 Nov 30;56(9-10):843-56

Zenger F et al (2004) Decreased glutathione in patients with anorexia nervosa. Risk factor for toxic liver injury? *Eur J Clin Nutr*. 2004 Feb;58(2):238-43

Zhong Z et al (2003) L-Glycine: a novel antiinflammatory, immunomodulatory, and cytoprotective agent. *Curr Opin Clin Nutr Metab Care*. 2003 Mar;6(2):229-40

References for Cysteine

Alexander B (1998) The role of nitric oxide in hepatic metabolism. *Nutrition*. 1998 Apr;14(4):376-90

Anderson ME, Luo JL (1998) Glutathione therapy: from prodrugs to genes. *Semin Liver Dis*. 1998;18(4):415-24

Bonkovsky HL (1997) Therapy of hepatitis C: other options. *Hepatology*. 1997 Sep;26(3 Suppl 1):143S-151S

Carmiel-Haggai M, Cederbaum AI, Nieto N (2003) Binge ethanol exposure increases liver injury in obese rats. *Gastroenterology*. 2003 Dec;125(6):1818-33

Carvalho M et al (2004) The toxicity of N-methyl-alpha-methyltyrosine to freshly isolated rat hepatocytes is prevented by ascorbic acid and N-acetylcysteine. *Toxicology*. 2004 Aug 5;200(2-3):193-203.

Coyle P et al (2002) Metallothionein: the multipurpose protein. *Cell Mol Life Sci*. 2002 Apr;59(4):627-47

Eisen JS et al (2004) N-acetylcysteine for the treatment of clove oil-induced fulminant hepatic failure. *J Toxicol Clin Toxicol*. 2004;42(1):89-92

Glantzounis GK et al (2004) Continuous infusion of N-acetylcysteine reduces liver warm ischaemia-reperfusion injury. *Br J Surg*. 2004 Oct;91(10):1330-9

Glazenburg EJ et al (1983) Effects of variation in the dietary supply of cysteine and methionine on liver concentration of glutathione and "active sulfate" (PAPS) and serum levels of sulfate, cysteine, methionine and taurine: relation to the metabolism of acetaminophen. *J Nutr*. 1983 Jul;113(7):1363-73

Jones AL (1998) Mechanism of action and value of N-acetylcysteine in the treatment of early and late acetaminophen poisoning: a critical review. *J Toxicol Clin Toxicol*. 1998;36(4):277-85

Kelly GS (1998) Clinical applications of N-acetylcysteine. *Altern Med Rev*. 1998 Apr;3(2):114-27

Krawinkel MB (2004) Parenteral nutrition-associated cholestasis--what do we know, what can we do? *Eur J Pediatr Surg*. 2004 Aug;14(4):230-4

Kretzschmar M (1996) Regulation of hepatic glutathione metabolism and its role in hepatotoxicity. *Exp Toxicol Pathol*. 1996 Jul;48(5):439-46

Lee TD et al (2004) Abnormal hepatic methionine and glutathione metabolism in patients with alcoholic hepatitis. *Alcohol Clin Exp Res*. 2004 Jan;28(1):173-81

Lieber CS (1997) Role of oxidative stress and antioxidant therapy in alcoholic and nonalcoholic liver diseases. *Adv Pharmacol*. 1997;38:601-28

Lieber CS (2002) S-adenosyl-L-methionine: its role in the treatment of liver disorders. *Am J Clin Nutr*. 2002 Nov;76(5):1183S-7S

Liu H et al (2004) Glutathione metabolism during aging and in Alzheimer disease. *Ann NY Acad Sci*. 2004 Jun;1019:346-9

Lu SC (1998) Regulation of hepatic glutathione synthesis. *Semin Liver Dis*. 1998;18(4):331-43

Majano PL et al (2004) N-Acetyl-cysteine modulates inducible nitric oxide synthase gene expression in human hepatocytes. *J Hepatol*. 2004 Apr;40(4):632-7

Mariotti F et al (2004) Acute ingestion of dietary proteins improves post-exercise liver glutathione in rats in a dose-dependent relationship with their cysteine content. *J Nutr*. 2004 Jan;134(1):128-31

Matsui H (2004) Sulfur-containing amino acids attenuate the development of liver fibrosis in rats through down-regulation of stellate cell activation. *J Hepatol*. 2004 Jun;40(6):917-25.

Patrick L (2003) Toxic metals and antioxidants: Part II. The role of antioxidants in arsenic and cadmium toxicity. *Altern Med Rev*. 2003 May;8(2):106-28

Rana SV, Allen T, Singh R (2002) Inevitable glutathione, then and now. *Indian J Exp Biol*. 2002 Jun;40(6):706-16

Ronchi VP et al (2004) The mouse liver content of carbonic anhydrase III and glutathione S-transferases A3 and P1 depend on dietary supply of methionine and cysteine. *Int J Biochem Cell Biol*. 2004 Oct;36(10):1993-2004

Sen CK (1999) Glutathione homeostasis in response to exercise training and nutritional supplements. *Mol Cell Biochem*. 1999 Jun;196(1-2):31-42

Soeters PB et al (2004) Amino acid adequacy in pathophysiological states. *J Nutr*. 2004 Jun;134(6 Suppl):1575S-1582S

Stipanuk MH (2004) Role of the liver in regulation of body cysteine and taurine levels: a brief review. *Neurochem Res*. 2004 Jan;29(1):105-10

Wu G et al (2004) Glutathione metabolism and its implications for health. *J Nutr*. 2004 Mar;134(3):489-92

Yamauchi A et al (1999) Liver immunity and glutathione. *Antioxid Redox Signal*. 1999 Summer;1(2):245-53

Yeh YY, Liu L (2001) Cholesterol-lowering effect of garlic extracts

References

- and organosulfur compounds: human and animal studies. *J Nutr.* 2001 Mar;131(3s):989S-93S
- Zafarullah M et al (2003) Molecular mechanisms of N-acetylcysteine actions. *Cell Mol Life Sci.* 2003 Jan;60(1):6-20
- Zenger F et al (2004) Decreased glutathione in patients with anorexia nervosa. Risk factor for toxic liver injury? *Eur J Clin Nutr.* 2004 Feb;58(2):238-43
- References for Glutamine**
- Buchman AL (2001) Glutamine: commercially essential or conditionally essential? A critical appraisal of the human data. *Am J Clin Nutr.* 2001 Jul;74(1):25-32
- Dadmarz M et al (1998) Effects of stress on amino acids and related compounds in various tissues of fasted rats. *Life Sci.* 1998;63(16):1485-91
- Dhar A, Kujath S, Van Way CW (2003) Glutamine administration during total parenteral nutrition protects liver adenosine nucleotides during and after subsequent hemorrhagic shock. *JPEN J Parenter Enteral Nutr.* 2003 Jul-Aug;27(4):246-51
- Kretzschmar M (1996) Regulation of hepatic glutathione metabolism and its role in hepatotoxicity. *Exp Toxicol Pathol.* 1996 Jul;48(5):439-46
- Lauterburg BH (2002) Analgesics and glutathione. *Am J Ther.* 2002 May-Jun;9(3):225-33
- Loguercio C et al (1999) Ethanol consumption, amino acid and glutathione blood levels in patients with and without chronic liver disease. *Alcohol Clin Exp Res.* 1999 Nov;23(11):1780-4
- Mates JM et al (2002) Glutamine and its relationship with intracellular redox status, oxidative stress and cell proliferation/death. *Int J Biochem Cell Biol.* 2002 May;34(5):439-58
- Newsholme P et al (2003) Glutamine and glutamate—their central role in cell metabolism and function. *Cell Biochem Funct.* 2003 Mar;21(1):1-9
- Nissim I (1999) Newer aspects of glutamine/glutamate metabolism: the role of acute pH changes. *Am J Physiol.* 1999 Oct;277(4 Pt 2):F493-7
- Olde Damink SW et al (2002) Interorgan ammonia metabolism in liver failure. *Neurochem Int.* 2002 Aug-Sep;41(2-3):177-88
- Rennie MJ et al (2001) Interaction between glutamine availability and metabolism of glycogen, tricarboxylic acid cycle intermediates and glutathione. *J Nutr.* 2001 Sep;131(9 Suppl):2488S-90S; discussion 2496S-7S
- Savilov PN (2004) The ammonia neutralization function of the liver in chronic active hepatitis. *Patol Fiziol Eksp Ter.* 2004 Jan-Mar;(1):24-6
- Soeters PB et al (2004) Amino acid adequacy in pathophysiological states. *J Nutr.* 2004 Jun;134(6 Suppl):1575S-1582S
- Watford M (2000) Glutamine and glutamate metabolism across the liver sinusoid. *J Nutr.* 2000 Apr;130(4S Suppl):983S-7S
- Watford M et al (2002) Hepatic glutamine metabolism. *Nutrition.* 2002 Apr;18(4):301-3
- Zenger F et al (2004) Decreased glutathione in patients with anorexia nervosa. Risk factor for toxic liver injury? *Eur J Clin Nutr.* 2004 Feb;58(2):238-43
- References for Lysine**
- Anaya-Prado R et al (2003) The attenuation of hemorrhage-induced liver injury by exogenous nitric oxide, L-arginine, and inhibition of inducible nitric oxide synthase. *J Invest Surg.* 2003 Sep-Oct;16(5):247-61
- Civitelli R et al (1992) Dietary L-lysine and calcium metabolism in humans. *Nutrition.* 1992 Nov-Dec;8(6):400-5
- Duncan AM, Ball RO, Pencharz PB (1996) Lysine requirement of adult males is not affected by decreasing dietary protein. *Am J Clin Nutr.* 1996 Nov;64(5):718-25
- Flodin NW (1997) The metabolic roles, pharmacology, and toxicology of lysine. *J Am Coll Nutr.* 1997 Feb;16(1):7-21
- Friedman M (1999) Chemistry, nutrition, and microbiology of D-amino acids. *J Agric Food Chem.* 1999 Sep;47(9):3457-79
- Hussain T et al (2004) Lysine fortification of wheat flour improves selected indices of the nutritional status of predominantly cereal-eating families in Pakistan. *Food Nutr Bull.* 2004 Jun;25(2):114-22
- Kurpad AV et al (2003) Lysine requirements of chronically undernourished adult Indian men, measured by a 24-h indicator amino acid oxidation and balance technique. *Am J Clin Nutr.* 2003 Jan;77(1):101-8
- Smriga M et al (2004) Lysine fortification reduces anxiety and lessens stress in family members in economically weak communities in Northwest Syria. *Proc Natl Acad Sci USA.* 2004 Jun 1;101(22):8285-8. Epub 2004 May 24
- Srivastava N et al (1980) Effect of lysine and threonine deficient and supplement rice diets on lipids of hepatic smooth and rough endoplasmic reticulum of rats. *Biochem Exp Biol.* 1980;16(4):443-4
- Thomas MR et al (1991) Lysine and protein metabolism in the young lactating woman. *Eur J Clin Nutr.* 1991 May;45(5):227-42
- Walsler M, Mitch WE, Abras E (1983) Supplements containing amino acids and keto acids in the treatment of chronic uremia. *Kidney Int Suppl.* 1983 Dec;16:S285-9
- Zello GA, Pencharz PB, Ball RO (1993) Dietary lysine requirement of young adult males determined by oxidation of L-[1-13C]phenylalanine. *Am J Physiol.* 1993 Apr;264(4 Pt 1):E677-85
- Zhao W et al (2004) Lysine-fortified wheat flour improves the nutritional and immunological status of wheat-eating families in northern China. *Food Nutr Bull.* 2004 Jun;25(2):123-9
- References for Taurine**
- Balkan J et al (2002) Improving effect of dietary taurine supplementation on the oxidative stress and lipid levels in the plasma, liver and aorta of rabbits fed on a high-cholesterol diet. *Biosci Biotechnol Biochem.* 2002 Aug;66(8):1755-8
- Chen Y, Li S, Zhang X (1999) Taurine inhibits deposition of extracellular matrix in experimental liver fibrosis in rats. *Zhonghua Gan Zang Bing Za Zhi.* 1999 Sep;7(3):165-7
- Chen YX (1993) Protective action of taurine on ischemia-reperfusion liver injury in rats and its mechanism. *Zhonghua Yi Xue Za Zhi.* 1993 May;73(5):276-9, 318-9
- Erman F et al (2004) Betaine or taurine administration prevents fibrosis and lipid peroxidation induced by rat liver by ethanol plus carbon tetrachloride intoxication. *Amino Acids.* 2004 Oct;27(2):199-205. Epub 2004 Aug 26
- Mashkova Nlu, Borovikova GV, Dokshina GA (1983) Effect of insulin and taurine on the potassium content of the perfused liver in irradiated rats. *Radiobiologia.* 1983 Mar-Apr;23(2):240-3
- Nakamura-Yamanaka Y, Tsuji K, Ichikawa T (1987) Effect of dietary taurine on cholesterol 7 alpha-hydroxylase activity in the liver of mice fed a lithogenic diet. *J Nutr Sci Vitaminol (Tokyo).* 1987 Jun;33(3):239-43
- Nakashima T et al (1996) Taurine in the liver. The function of taurine conjugated with bile acids. *Adv Exp Med Biol.* 1996;403:85-92
- Obinata K et al (1996) Effect of taurine on the fatty liver of children with simple obesity. *Adv Exp Med Biol.* 1996;403:607-13
- Park T, Lee K (1998) Dietary taurine supplementation reduces plasma and liver cholesterol and triglyceride levels in rats fed a high-cholesterol or a cholesterol-free diet. *Adv Exp Med Biol.* 1998;442:319-25
- Pierre Y, Chatagner F (1981) A possible physiological role of taurine in the adult female rat liver. *Reprod Nutr Dev.* 1981;21(4):555-60
- Refik Mas M et al (2004) The effect of taurine treatment on oxidative stress in experimental liver fibrosis. *Hepatol Res.* 2004 Apr;28(4):207-215
- Salas Martinez J et al (1994) Parenteral nutrition-associated cholestasis: the effect of diets free of taurine and its metabolic precursors on bile composition and liver structure. *Nutr Hosp.* 1994 Jan-Feb;9(1):44-53
- Scanlan N (2001) Compromised hepatic detoxification in companion animals and its correction via nutritional supplementation and modified fasting. *Altern Med Rev.* 2001 Sep;6 Suppl:S24-37
- Soeters PB et al (2004) Amino acid adequacy in pathophysiological states. *J Nutr.* 2004 Jun;134(6 Suppl):1575S-1582S
- Waterfield CJ et al (1993) Reduction of liver taurine in rats by beta-alanine treatment increases carbon tetrachloride toxicity. *Toxicology.* 1993 Jan 29;77(1-2):7-20
- Waterfield CJ et al (1993a) The correlation between urinary and liver taurine levels and between pre-dose urinary taurine and liver damage. *Toxicology.* 1993 Jan 29;77(1-2):1-5
- Waterfield CJ et al (1993b) Investigations into the effects of various hepatotoxic compounds on urinary and liver taurine levels in rats. *Arch Toxicol.* 1993;67(4):244-54
- Waterfield CJ et al (1993c) Effect of various non-hepatotoxic compounds on urinary and liver taurine levels in rats. *Arch Toxicol.* 1993;67(8):538-46
- References for Carnitine**
- Allard JP (2002) Other disease associations with non-alcoholic fatty liver disease (NAFLD). *Best Pract Res Clin Gastroenterol.* 2002 Oct;16(5):783-95
- Bach AC (1982) Carnitine in human nutrition. *Z Ernährungswiss.* 1982 Dec;21(4):257-65
- Brivet M et al (1999) Defects in activation and transport of fatty acids. *J Inherit Metab Dis.* 1999 Jun;22(4):428-41
- Bykov I, Jarvelainen H, Lindros K (2003) L-carnitine alleviates alcohol-induced liver damage in rats: role of tumour necrosis factor-alpha. *Alcohol Alcohol.* 2003 Sep-Oct;38(5):400-6
- Cha YS, Kim HY, Daily JW 3rd. (2003) Exercise-trained but not untrained rats maintain free carnitine reserves during acute exercise. *Asia Pac J Clin Nutr.* 2003;12(1):120-6
- Chitturi S, George J (2002) Hepatotoxicity of commonly used drugs: nonsteroidal anti-inflammatory drugs, antihypertensives, antidiabetic agents, anticonvulsants, lipid-lowering agents, psychotropic drugs. *Semin Liver Dis.* 2002;22(2):169-83
- Czech E, Olszowy Z, Nowicka J (2004) The influence of L-carnitine on methanol biotransformation in rats. *Exp Toxicol Pathol.* 2004 Mar;55(5):367-77
- Demirdag K et al (2004) Role of L-carnitine in the prevention of acute liver damage induced by carbon tetrachloride in rats. *J Gastroenterol Hepatol.* 2004 Mar;19(3):333-8
- Evangelidou A, Vlassopoulos D (2003) Carnitine metabolism and deficit—when supplementation is necessary? *Curr Pharm Biotechnol.* 2003 Jun;4(3):211-9
- Ibrahim VH et al (2003) Effects of carnitine and taurine on fatty acid metabolism and lipid accumulation in the liver of cats during weight gain and weight loss. *Am J Vet Res.* 2003 Oct;64(10):1265-77
- Krahenbuhl S (1996) Carnitine metabolism in chronic liver disease. *Life Sci.* 1996;59(19):1579-99
- Lapinski TW, Grzeszczuk A (2003) The impact of carnitine on serum ammonia concentration and lipid metabolism in patients with alcoholic liver cirrhosis. *Pol Merkuriusz Lek.* 2003 Jul;15(85):38-41
- Li XX et al (2002) Hyperammonemia in carnitine-deficient adult JVS mice used by starvation. *Metab Brain Dis.* 2002 Dec;17(4):359-66
- Malaguarnera M et al (2003) L-Carnitine in the treatment of mild or moderate hepatic encephalopathy. *Dig Dis.* 2003;21(3):271-5
- Neri S et al (2003) L-carnitine decreases severity and type of fatigue induced by interferon-alpha in the treatment of patients with hepatitis C. *Neuropsychobiology.* 2003;47(2):94-7
- Perez-Carreras M et al (2003) Defective hepatic mitochondrial respiratory chain in patients with nonalcoholic steatohepatitis. *Hepatology.* 2003 Oct;38(4):999-1007
- Schmid A, Schmid H (1996) Possibilities and limitations of drugs to protect the liver *Tierarztl Prax.* 1996 Dec;24(6):537-41
- Selimoglu MA, Yagci RV (2004) Plasma and liver carnitine levels of children with chronic hepatitis B. *J Clin Gastroenterol.* 2004 Feb;38(2):130-3
- Spaniol M et al (2003) Mechanisms of liver steatosis in rats with systemic carnitine deficiency due to treatment with trimethylhydraziniumpropionate. *J Lipid Res.* 2003 Jan;44(1):144-53
- Spriet LL, Peters SJ (1998) Influence of diet on the metabolic responses to exercise. *Proc Nutr Soc.* 1998 Feb;57(1):25-33
- Steiber A, Kerner J, Hoppel CL (2004) Carnitine: a nutritional, biosynthetic, and functional perspective. *Mol Aspects Med.* 2004 Oct-Dec;25(5-6):455-73
- Szilagy M (1998) L-carnitine as essential methylated compound in animal metabolism. An overview. *Acta Biol Hung.* 1998;49(2-4):209-18
- Tein I (2003) Carnitine transport: pathophysiology and metabolism of known molecular defects. *J Inherit Metab Dis.* 2003;26(2-3):147-69
- Washington L, Cook GA, Mansbach CM (2003) Inhibition of carnitine palmitoyltransferase in the rat small intestine reduces export of triacylglycerol into the lymph. *J Lipid Res.* 2003 Jul;44(7):1395-403. Epub 2003 Apr 16
- References for Glutamic acid**
- Brosnan JT (2000) Glutamate, at the interface between amino acid and carbohydrate metabolism. *J Nutr.* 2000 Apr;130(4S Suppl):988S-90S
- Butterworth RF (2002b) Glutamate transporters in hyperammonemia. *Neurochem Int.* 2002 Aug-Sep;41(2-3):81-5
- Chambon-Savanovitch C et al (1999) Can a glutamate-enriched diet counteract glutamine depletion in endotoxemic rats? *J Nutr Biochem.* 1999 Jun;10(6):331-7
- Dadmarz M et al (1998) Effects of stress on amino acids and related compounds in various tissues of fasted rats. *Life Sci.* 1998;63(16):1485-91
- Jones EA (2000) Pathogenesis of hepatic encephalopathy. *Clin Liver Dis.* 2000 May;4(2):467-85
- Kelly A, Stanley CA (2001) Disorders of glutamate metabolism. *Ment Retard Dev Disabil Res Rev.* 2001;7(4):287-95
- Kretzschmar M (1996) Regulation of hepatic glutathione metabolism and its role in hepatotoxicity. *Exp Toxicol Pathol.* 1996 Jul;48(5):439-46
- Lauterburg BH (2002) Analgesics and glutathione. *Am J Ther.* 2002 May-Jun;9(3):225-33
- Loguercio C et al (1999) Ethanol consumption, amino acid and glutathione blood levels in patients with and without chronic liver disease. *Alcohol Clin Exp Res.* 1999 Nov;23(11):1780-4
- Newsholme P et al (2003) Glutamine and glutamate—their central role in cell metabolism and function. *Cell Biochem Funct.* 2003 Mar;21(1):1-9
- Nissim I (1999) Newer aspects of glutamine/glutamate metabolism: the role of acute pH changes. *Am J Physiol.* 1999 Oct;277(4 Pt 2):F493-7
- Nissim I et al (2003) Role of the glutamate dehydrogenase reaction in furnishing aspartate nitrogen for urea synthesis: studies in perfused rat liver with 15N. *Biochem J.* 2003 Nov 15;376(Pt 1):179-88
- Olde Damink SW et al (2002) Interorgan ammonia metabolism in liver failure. *Neurochem Int.* 2002 Aug-Sep;41(2-3):177-88
- Rennie MJ et al (2001) Interaction between glutamine availability and metabolism of glycogen, tricarboxylic acid cycle intermediates and glutathione. *J Nutr.* 2001 Sep;131(9 Suppl):2488S-90S; discussion 2496S-7S
- Yang D, Brunengraber H (2000) Glutamate, a window on liver intermediary metabolism. *J Nutr.* 2000 Apr;130(4S Suppl):991S-4S

References

References for Inositol

- Albi E, Viola Magni MP (2004) The role of intranuclear lipids. *Biol Cell*. 2004 Oct;96(8):657-67
- Bokkala S, Rubin E, Joseph SK (1999) Effect of chronic ethanol exposure on inositol triphosphate receptors in WB rat liver epithelial cells. *Alcohol Clin Exp Res*. 1999 Dec;23(12):1875-83
- Burgess JW et al (2003) Phosphatidylinositol promotes cholesterol transport and excretion. *J Lipid Res*. 2003 Jul;44(7):1355-63. Epub 2003 Apr 16
- Dufour JF et al (1999) Expression of inositol 1,4,5-trisphosphate receptor isoforms in rat cirrhosis. *Hepatology*. 1999 Oct;30(4):1018-26
- Fox CH, Eberl M (2002) Phytic acid (IP6), novel broad spectrum anti-neoplastic agent: a systematic review. *Complement Ther Med*. 2002 Dec;10(4):229-34
- Grases F et al (2001) Phytate levels in diverse rat tissues: influence of dietary phytate. *Br J Nutr*. 2001 Aug;86(2):225-31
- Holub BJ (1982) The nutritional significance, metabolism, and function of myo-inositol and phosphatidylinositol in health and disease. *Adv Nutr Res*. 1982;4:107-41
- Holub BJ (1986) Metabolism and function of myo-inositol and inositol phospholipids. *Annu Rev Nutr*. 1986;6:563-97
- Jarivalla RJ (2001) Rice-bran products: phytonutrients with potential applications in preventive and clinical medicine. *Drugs Exp Clin Res*. 2001;27(1):17-26
- Katayama T (1999) Hypolipidemic action of phytic acid (IP6): prevention of fatty liver. *Anticancer Res*. 1999 Sep-Oct;19(5A):3695-8
- Nishino H et al (1999) Suppression of lung and liver carcinogenesis in mice by oral administration of myo-inositol. *Anticancer Res*. 1999 Sep-Oct;19(5A):3663-4
- Okazaki Y, Katayama T (2003) Effects of dietary carbohydrate and myo-inositol on metabolic changes in rats fed 1,1,1-trichloro-2,2-bis (p-chlorophenyl) ethane (DDT). *J Nutr Biochem*. 2003 Feb;14(2):81-9
- Pusl T, Nathanson MH (2004) The role of inositol 1,4,5-trisphosphate receptors in the regulation of bile secretion in health and disease. *Biochem Biophys Res Commun*. 2004 Oct 1;322(4):1318-25
- Rakhmatullaev FKh, Khakimov ZZ (2001) Phytin efficacy in correction of the impaired detoxication liver function in the long-term compression syndrome in rats in prepubertal period. *Ekspl Klin Farmakol*. 2001 Jan-Feb;64(1):71-3
- Santi P et al (2003) Inositol-specific phospholipase C in low and fast proliferating hepatoma cell lines. *Int J Oncol*. 2003 May;22(5):1147-53
- Sun TH et al (2002) Both myo-inositol to chiro-inositol epimerase activities and chiro-
- Sun TH et al (2002) Both myo-inositol to chiro-inositol epimerase activities and chiro-inositol to myo-inositol ratios are decreased in tissues of GK type 2 diabetic rats compared to Wistar controls. *Biochem Biophys Res Commun*. 2002 May 10;293(3):1092-8
- Skudelski T, Kandulski K (1999) The effect of myo-inositol on ethanol-induced metabolic changes and insulin concentration in the rat. *Arch Physiol Biochem*. 1999 Oct;107(4):334-7

References for Lecithin

- Aleynik MK, Lieber CS (2001) Dilinoleoylphosphatidylcholine decreases ethanol-induced cytochrome P450E1. *Biochem Biophys Res Commun*. 2001 Nov 9;288(4):1047-51
- Aleynik SI, Lieber CS (2003) Polyenyolphosphatidylcholine corrects the alcohol-induced hepatic oxidative stress by restoring s-adenosylmethionine. *Alcohol Alcohol*. 2003 May-Jun;38(3):208-12
- Baraona E et al (2002) Ethanol consumption increases nitric oxide production in rats, and its peroxynitrite-mediated toxicity is attenuated by polyenyolphosphatidylcholine. *Alcohol Clin Exp Res*. 2002 Jun;26(6):883-9
- Cabre E, Gassull MA (1995) Nutritional support in liver disease. *Eur J Gastroenterol Hepatol*. 1995 Jun;7(6):528-32
- Carrasco MP et al (2002) Comparative study of the effects of short- and long-term ethanol treatment and alcohol withdrawal on phospholipid biosynthesis in rat hepatocytes. *Comp Biochem Physiol B Biochem Mol Biol*. 2002 Mar;131(3):491-7
- Chanussot F, Benkoel L (2003) Prevention by dietary (n-6) polyunsaturated phosphatidylcholines of intrahepatic cholestasis induced by cyclosporine A in animals. *Life Sci*. 2003 Jun 13;73(4):381-92.
- Demyda EP (2003) Cholesecretory function of the liver and correction of its disorders in patients with cholesterosis of the gallbladder. *Lik Sprava*. 2003 Apr-Jun;3(4):68-72
- Halsted CH (2004) Nutrition and alcoholic liver disease. *Semin Liver Dis*. 2004 Aug;24(3):289-304
- Karaman A et al (2003) Protective effect of polyunsaturated phosphatidylcholine on liver damage induced by biliary obstruction in rats. *J Pediatr Surg*. 2003 Sep;38(9):1341-7
- Kasbo J et al (2003) Phosphatidylcholine-enriched diet prevents gallstone formation in mice susceptible to cholelithiasis. *J Lipid Res*. 2003 Dec;44(12):2297-303. Epub 2003 Jul 01
- Lata J et al (2001) Protective effect of essential phospholipids on liver injury due to total parenteral nutrition. *Vnitr Lek*. 2001 Sep;47(9):599-603

- LeBlanc MJ et al (2003) Effects of dietary soybean lecithin on plasma lipid transport and hepatic cholesterol metabolism in rats. *J Nutr Biochem*. 2003 Jan;14(1):40-8
- Lieber CS (1993) Biochemical factors in alcoholic liver disease. *Semin Liver Dis*. 1993 May;13(2):136-53
- Lieber CS (1999) Prevention and treatment of liver fibrosis based on pathogenesis. *Alcohol Clin Exp Res*. 1999 May;23(5):944-9
- Lieber CS et al (2003) I. Veterans Affairs Cooperative Study of polyenyolphosphatidylcholine in alcoholic liver disease: effects on drinking behavior by nurse/physician teams. *Alcohol Clin Exp Res*. 2003 Nov;27(11):1757-64
- Lieber CS et al (2003b) II. Veterans Affairs Cooperative Study of polyenyolphosphatidylcholine in alcoholic liver disease. *Alcohol Clin Exp Res*. 2003 Nov;27(11):1765-72
- Noga AA, Vance DE (2003) Insights into the requirement of phosphatidylcholine synthesis for liver function in mice. *J Lipid Res*. 2003 Oct;44(10):1998-2005. Epub 2003 Jul 01
- Reo NV, Adinezhadeh M, Foy BD (2002) Kinetic analyses of liver phosphatidylcholine and phosphatidylethanolamine biosynthesis using ¹³C NMR spectroscopy. *Biochim Biophys Acta*. 2002 Feb 28;1580(2-3):171-88
- Turecky L et al (2003) Plasma lipid parameters in patients with alcoholic fatty liver after treatment with essential phospholipids. *Bratisl Lek Listy*. 2003;104(7-8):227-31
- Vance JE, Vance DE (1985) The role of phosphatidylcholine biosynthesis in the secretion of lipoproteins from hepatocytes. *Can J Biochem Cell Biol*. 1985 Aug;63(8):870-81

References for Potassium

- Berrino F (2002) Western diet and Alzheimer's disease. *Epidemiol Prev*. 2002 May-Jun;26(3):107-15
- Chukhlova EA, Kudzhina Llu, Evtdienko IuV (1982) Effect of fasting on the content and transport of potassium ions in rat liver mitochondria. *Ukr Biokhim Zh*. 1982 Mar-Apr;54(2):190-3
- Decaux G et al (1988) Potassium homeostasis in liver cirrhosis. *Arch Intern Med*. 1988 Mar;148(3):547-8
- Kabakov Alu, Sarbash VI, Filimonov DA (1989) Potassium transport in ischemia and anoxia of the perfused liver. *Biofizika*. 1989 Mar-Apr;34(2):295-6
- Mashkova Nlu, Borovikova GV, Dokshina GA (1983) Effect of insulin and taurine on the potassium content of the perfused liver in irradiated rats. *Radiobiologia*. 1983 Mar-Apr;23(2):240-3
- Nietsch HH et al (2000) Activation of potassium and chloride channels by tumor necrosis factor alpha. Role in liver cell death. *J Biol Chem*. 2000 Jul 7;275(27):20556-61
- Shishiba Y et al (1980) Potassium deficiency enhances the effect of thyroid hormone on NaK-ATPase in liver and kidney. *Endocrinol Jpn*. 1980 Jun;27(3):329-36

References for Magnesium

- Berrino F (2002) Western diet and Alzheimer's disease. *Epidemiol Prev*. 2002 May-Jun;26(3):107-15
- Bojanovic K, Skarbek-Galamon C (1977) Disturbances of fat, carbohydrate, magnesium and protein balance in liver diseases. *Acta Hepatogastroenterol (Stuttg)*. 1977 Jun;24(3):155-61
- Chiba T et al (1987) Hypocalcemic crisis in alcoholic fatty liver: transient hypoparathyroidism due to magnesium deficiency. *Am J Gastroenterol*. 1987 Oct;82(10):1084-7
- Cohen L (1985) Magnesium and liver cirrhosis: a hypothesis. *Magnesium*. 1985;4(1):1-4
- Koivisto M et al (2002) Magnesium depletion in chronic terminal liver cirrhosis. *Clin Transplant*. 2002 Oct;16(5):325-8
- Oralewska B et al (1996) Disorders of magnesium homeostasis in the course of liver disease in children. *Magn Res*. 1996 Jun;9(2):125-8
- Patrick L (2002) Nonalcoholic fatty liver disease: relationship to insulin sensitivity and oxidative stress. Treatment approaches using vitamin E, magnesium, and betaine. *Altern Med Rev*. 2002 Aug;7(4):276-91
- Rayssiguier Y et al (1985) Influence of magnesium deficiency on liver collagen after carbon tetrachloride or ethanol administration to rats. *J Nutr*. 1985 Dec;115(12):1656-62
- Reis MA et al (2000) Magnesium deficiency modulates the insulin signaling pathway in liver but not muscle of rats. *J Nutr*. 2000 Feb;130(2):133-8
- Rocchi E et al (1994) Zinc and magnesium in liver cirrhosis. *Eur J Clin Invest*. 1994 Mar;24(3):149-55
- Rocchi E et al (1994b) Zinc and magnesium in liver cirrhosis. *Eur J Clin Invest*. 1994 Mar;24(3):149-55
- Zozycka Z, Wutzn J, Lewicki Z (1978) Fat-storing cells in the liver of rats kept on a low-magnesium diet. *Exp Pathol (Jena)*. 1978;15(6):335-40

References for Aspartic acid

- Butterworth RF (2003) Role of circulating neurotoxins in the pathogenesis of hepatic encephalopathy: potential for improvement following their removal by liver assist devices. *Liver Int*. 2003;23 Suppl 3:5-9
- Butterworth RF (2002) Pathophysiology of hepatic encephalopathy: a new look at ammonia. *Metab Brain Dis*. 2002 Dec;17(4):221-7
- Gebhardt R et al (1997) Treatment of cirrhotic rats with L-ornithine-

- L-aspartate enhances urea synthesis and lowers serum ammonia levels. *J Pharmacol Exp Ther*. 1997 Oct;283(1):1-6
- Muting D, Reikowski J (1983) Hepatic coma. Principles of pathogenesis and treatment. 2: Treatment, prognosis. *Fortschr Med*. 1983 Oct 20;101(39):1766-73
- Nissim I et al (2003) Role of the glutamate dehydrogenase reaction in furnishing aspartate nitrogen for urea synthesis: studies in perfused rat liver with ¹⁵N. *Biochem J*. 2003 Nov 15;376(Pt 1):179-88
- Stauch S et al (1998) Oral L-ornithine-L-aspartate therapy of chronic hepatic encephalopathy: results of a placebo-controlled double-blind study. *J Hepatol*. 1998 May;28(5):856-64
- Tandon BN (2003) Hepatic encephalopathy syndromes. *Indian J Gastroenterol*. 2003 Dec;22 Suppl 2:S4-6
- Weijjs PJ et al (1996) Conversion of [¹⁵N]ammonia into urea and amino acids in humans and the effect of nutritional status. *Br J Nutr*. 1996 Oct;76(4):491-9
- Yassuda FP et al (2003) The urea cycle in the liver of arthritic rats. *Mol Cell Biochem*. 2003 Jan;243(1-2):97-106

References for Zinc

- Ajayi OB, Oduyiga A (2004) Effect of low-zinc status and essential fatty acids deficiency on the activities of aspartate aminotransferase and alanine aminotransferase in liver and serum of albino rats. *Nahrung*. 2004 Apr;48(2):88-90
- Al-Bader AA et al (1997) Effect of dietary selenium, zinc and allopurinol supplements on plasma and tissue manganese levels in rats with thioacetamide [correction of thiocetamide]-induced liver cirrhosis. *Mol Cell Biochem*. 1997 Aug;173(1-2):121-5
- Baltaci AK et al (2003) Effects of zinc deficiency and supplementation on the glycogen contents of liver and plasma lactate and leptin levels of rats performing acute exercise. *Biol Trace Elem Res*. 2003 Winter;96(1-3):227-36
- Barry M, Keeling PW, Feely J (1990) Tissue zinc status and drug elimination in patients with chronic liver disease. *Clin Sci (Lond)*. 1990 Jun;78(6):547-9
- Capocaccia L et al (1991) Zinc and other trace elements in liver cirrhosis. *Ital J Gastroenterol*. 1991 Jul-Aug;23(6):386-91
- Chen JF, Shi QJ, Zheng SY (2001) Experimental research of protective and therapeutic effects of zinc and vitamin E on mouse liver radiational damage. *Hunan Yi Ke Da Xue Xue Bao*. 2001 Jun 28;26(3):207-10
- Dashti HM et al (1997) Zinc and liver cirrhosis: biochemical and histopathologic assessment. *Nutrition*. 1997 Mar;13(3):206-12
- Dhawan D, Goel A (1995) Further evidence for zinc as a hepatoprotective agent in rat liver toxicity. *Exp Mol Pathol*. 1995 Oct;63(2):110-7
- Goel A, Chauhan DP, Dhawan DK (2000) Protective effects of zinc in chlorpyrifos induced hepatotoxicity: a biochemical and trace elemental study. *Biol Trace Elem Res*. 2000 May;74(2):171-83
- Goel A, Dhawan DK (2001) Zinc supplementation prevents liver injury in chlorpyrifos-treated rats. *Biol Trace Elem Res*. 2001 Summer;82(1-3):185-200
- Grungreiff K, Reinhold D (2005) Liver cirrhosis and "liver" diabetes mellitus are linked by zinc deficiency. *Med Hypotheses*. 2005;64(2):316-7
- Gur G et al (1998) Determination of hepatic zinc content in chronic liver disease due to hepatitis B virus. *Hepatogastroenterology*. 1998 Mar-Apr;45(20):472-6
- Hsu PC, Guo YL (2002) Antioxidant nutrients and lead toxicity. *Toxicology*. 2002 Oct 30;180(1):33-44
- Kawase M et al (2004) Liver protection by bis(maltolato)zinc(III) complex. *Exp Anim*. 2004 Jan;53(1):1-9
- Marchetti P et al (1998) Zinc deficiency in liver cirrhosis: a curiosity or a problem? *Ann Ital Med Int*. 1998 Jul-Sep;13(3):157-62
- Nomura F, Takekoshi K (1994) Zinc and selenium metabolism in liver cirrhosis. *Nippon Rinsho*. 1994 Jan;52(1):165-9
- Patrick L (2003) Toxic metals and antioxidants: Part II. The role of antioxidants in arsenic and cadmium toxicity. *Altern Med Rev*. 2003 May;8(2):106-28
- Poschl G, Seitz HK (2004) Alcohol and cancer. *Alcohol*. 2004 May-Jun;39(3):155-65
- Riggio O et al (1992) Zinc supplementation reduces blood ammonia and increases liver ornithine transcarbamylase activity in experimental cirrhosis. *Hepatology*. 1992 Sep;16(3):785-9
- Rocchi E et al (1994b) Zinc and magnesium in liver cirrhosis. *Eur J Clin Invest*. 1994 Mar;24(3):149-55
- Rodriguez-Moreno F et al (1997) Zinc, copper, manganese, and iron in chronic alcoholic liver disease. *Alcohol*. 1997 Jan-Feb;14(1):39-44
- Song YM, Chen MD (2003) Zinc supplementation attenuates thioacetamide-induced liver injury and hyperglycemia in mice. *Biol Trace Elem Res*. 2003 May;92(2):173-80
- Torra M et al (1995) Cadmium and zinc relationships in the liver and kidney of humans exposed to environmental cadmium. *Sci Total Environ*. 1995 Aug 18;170(1-2):53-7
- Tsocheva-Gaitandjeva NT, Gabrashanska MP, Tepavitcharova S (2002) Trace element levels in the liver of rats with acute and chronic fascioliasis and after treatment with zinc-copper hydroxochloride mixed crystals. *J Helminthol*. 2002 Mar;76(1):87-90

References

- Yoshida Y et al (2001) Effects of zinc deficiency/zinc supplementation on ammonia metabolism in patients with decompensated liver cirrhosis. *Acta Med Okayama*. 2001 Dec;55(6):349-55
- Zhou Z et al (2002) Metallothionein-independent zinc protection from alcoholic liver injury. *Am J Pathol*. 2002 Jun;160(6):2267-74
- References for Selenium**
- Aboutwerat A et al (2003) Oxidant stress is a significant feature of primary biliary cirrhosis. *Biochim Biophys Acta*. 2003 Mar 20;1637(2):142-50
- Berghelm I et al (2003) Nutritional deficiencies in German middle-class male alcohol consumers: relation to dietary intake and severity of liver disease. *Eur J Clin Nutr*. 2003 Mar;57(3):431-8
- Beytut E, Aksakal M (2003) Effects of dietary vitamin E and selenium on antioxidative defense mechanisms in the liver of rats treated with high doses of glucocorticoid. *Biol Trace Elem Res*. 2003 Mar;91(3):231-41
- Bjelakovic G et al (2004) Antioxidant supplements for prevention of gastrointestinal cancers: a systematic review and meta-analysis. *Lancet*. 2004 Oct 2;364(9441):1219-28
- Brown KM, Arthur JR (2001) Selenium, selenoproteins and human health: a review. *Public Health Nutr*. 2001 Apr;4(2B):593-9
- Burk RF, Hill KE, Motley AK (2003) Selenoprotein metabolism and function: evidence for more than one function for selenoprotein P. *J Nutr*. 2003 May;133(5 Suppl 1):1517S-20S
- Carlson BA et al (2004) Specific excision of the selenocysteine tRNA(Ser)Sec (Trsp) gene in mouse liver demonstrates an essential role of selenoproteins in liver function. *J Biol Chem*. 2004 Feb 27;279(9):8011-7. Epub 2003 Dec 04
- Czuczajko J et al (2003) Selenium, glutathione and glutathione peroxidases in blood of patients with chronic liver diseases. *Acta Biochim Pol*. 2003;50(4):1147-54
- El-Demerdash FM (2004) Antioxidant effect of vitamin E and selenium on lipid peroxidation, enzyme activities and biochemical parameters in rats exposed to aluminium. *J Trace Elem Med Biol*. 2004;18(1):113-21
- Erisir M et al (2003) Effects of dietary vitamin E and selenium on arginase activity in the liver, kidneys, and heart of rats treated with high doses of glucocorticoid. *Cell Biochem Funct*. 2003 Dec;21(4):331-5
- Gehring MM et al (2003) An investigation into the effect of selenium supplementation on microcystin hepatotoxicity. *Toxicol*. 2003 Mar;41(4):451-8
- He YT et al (2004) Therapeutic effects and molecular mechanisms of anti-fibrosis herbs and selenium on rats with hepatic fibrosis. *World J Gastroenterol*. 2004 Mar 1;10(5):703-6
- Hsu PC, Guo YL (2002) Antioxidant nutrients and lead toxicity. *Toxicology*. 2002 Oct 30;180(1):33-44
- Ikemoto T et al (2004) Detoxification mechanism of heavy metals in marine mammals and seabirds: interaction of selenium with mercury, silver, copper, zinc, and cadmium in liver. *Arch Environ Contam Toxicol*. 2004 Oct;47(3):402-13.
- Jablonska-Kaszewska I et al (2003) Serum selenium levels in alcoholic liver disease. *Med Sci Monit*. 2003 Aug;9 Suppl 3:15-18
- Kim HY, Wallig MA, Picciano MF (2003) Inorganic selenite supplementation and protection against hyperoxic injury in neonates. *J Nutr Sci Vitaminol (Tokyo)*. 2003 Oct;49(5):350-6.
- Konovalova GG et al (2003) A complex of antioxidant vitamins effectively inhibits free-radical oxidation of LDL phospholipids in blood plasma and membrane structures of the liver and myocardium. *Bull Exp Biol Med*. 2003 Feb;135(2):143-6
- Lindsay LA et al (2004) Effect of daily cod liver oil and a multivitamin-mineral supplement with selenium on upper respiratory tract pediatric visits by young, inner-city, Latino children: randomized pediatric sites. *Ann Otol Rhinol Laryngol*. 2004 Nov;113(11):891-901
- Loguerio C, Federico A (2003) Oxidative stress in viral and alcoholic hepatitis. *Free Radic Biol Med*. 2003 Jan 1;34(1):1-10
- Lyons GH, Stangoulis JC, Graham RD (2004) Exploiting micronutrient interaction to optimize biofortification programs: the case for inclusion of selenium and iodine in the HarvestPlus program. *Nutr Rev*. 2004 Jun;62(6 Pt 1):247-52
- Matsumoto K et al (2002) Evaluation of oxidative damage in the liver of selenium-deficient rats. *Redox Rep*. 2002;7(5):351-4
- Navarro-Alarcon M, Lopez-Martinez MC (2000) Essentiality of selenium in the human body: relationship with different diseases. *Sci Total Environ*. 2000 Apr 17;249(1-3):347-71
- Naziroglu M, Karaoglu A, Aksoy AO (2004) Selenium and high dose vitamin E administration protects cisplatin-induced oxidative damage to renal, liver and lens tissues in rats. *Toxicology*. 2004 Feb 15;195(2-3):221-30
- Olsson U (1985) Selenium deficiency and detoxication functions in the rat: effect of chronic dietary cadmium. *Drug Nutr Interact*. 1985;3(3):129-40
- Ozdil S et al (2004) Protective effects of ascorbic acid, dl-alpha-tocopherol acetate, and sodium selenate on ethanol-induced liver damage of rats. *Biol Trace Elem Res*. 2004 Feb;97(2):149-62
- Park ES et al (2003) Protective effects of N-acetylcysteine and selenium against doxorubicin toxicity in rats. *J Vet Sci*. 2003 Aug;4(2):129-36
- Patrick L (2003) Toxic metals and antioxidants: Part II. The role of antioxidants in arsenic and cadmium toxicity. *Altern Med Rev*. 2003 May;8(2):106-28
- Poschl G, Seitz HK (2004) Alcohol and cancer. *Alcohol*. 2004 May-Jun;39(3):155-65
- Reddy CC, Massaro EJ (1983) Biochemistry of selenium: a brief overview. *Fundam Appl Toxicol*. 1983 Sep-Oct;3(5):431-6
- Reid ME et al (2004) A report of high-dose selenium supplementation: response and toxicities. *J Trace Elem Med Biol*. 2004;18(1):69-74
- Scanlan N (2001) Compromised hepatic detoxification in companion animals and its correction via nutritional supplementation and modified fasting. *Altern Med Rev*. 2001 Sep;6 Suppl:S24-37
- Schafer K et al (2004) Effects of selenium deficiency on fatty acid metabolism in rats fed fish oil-enriched diets. *Trace Elem Med Biol*. 2004;18(1):89-97
- Schwedhelm E et al (2003) Clinical pharmacokinetics of antioxidants and their impact on systemic oxidative stress. *Clin Pharmacokinet*. 2003;42(5):437-59
- Sivaram AG, Suresh MV, Indira M (2003) Combined effect of ascorbic acid and selenium supplementation on alcohol-induced oxidative stress in guinea pigs. *Comp Biochem Physiol C Toxicol Pharmacol*. 2003 Mar;134(3):397-401
- Soares JC, Folmer V, Rocha JB (2003) Influence of dietary selenium supplementation and exercise on thiol-containing enzymes in mice. *Nutrition*. 2003 Jul-Aug;19(7-8):627-32
- Soriano-Garcia M (2004) Organoselenium compounds as potential therapeutic and chemopreventive agents: a review. *Curr Med Chem*. 2004 Jun;11(12):1657-69
- Thirunavukkarasu C, Sakthisekaran D (2003) Sodium selenite, dietary micronutrient, prevents the lymphocyte DNA damage induced by N-nitrosodiethylamine and phenobarbital promoted experimental hepatocarcinogenesis. *J Cell Biochem*. 2003 Feb 15;88(3):578-88
- Venardos K et al (2004) Effects of dietary selenium on glutathione peroxidase and thioredoxin reductase activity and recovery from cardiac ischemia-reperfusion. *J Trace Elem Med Biol*. 2004;18(1):81-88
- Weiller M et al (2004) Toxicity of nutritionally available selenium compounds in primary and transformed hepatocytes. *Toxicology*. 2004 Sep 1;201(1-3):21-30
- Yu RA, Chen XM (2004) Effects of selenium on rat hepatocellular DNA damage induced by cadmium in vitro. *Zhonghua Yu Fang Yi Xue Za Zhi*. 2004 Jan;38(1):29-32
- References for Folic acid**
- Berrino F (2002) Western diet and Alzheimer's disease. *Epidemiol Prev*. 2002 May-Jun;26(3):107-15
- Branda RF et al (2002) Diet modulates the toxicity of cancer chemotherapy in rats. *J Lab Clin Med*. 2002 Nov;140(5):358-68
- Brunaud L et al (2003) Effects of vitamin B12 and folate deficiencies on DNA methylation and carcinogenesis in rat liver. *Clin Chem Lab Med*. 2003 Aug;41(8):1012-9
- Choy PC et al (2004) Lipids and atherosclerosis. *Biochem Cell Biol*. 2004 Feb;82(1):212-24
- Donnelly JG (2001) Folic acid. *Crit Rev Clin Lab Sci*. 2001 Jun;38(3):183-223.
- Ghandour H et al (2002) Folate status and age affect the accumulation of L-isoaspartyl residues in rat liver proteins. *J Nutr*. 2002 Jun;132(6):1357-60
- Halsted CH (2004) Nutrition and alcoholic liver disease. *Semin Liver Dis*. 2004 Aug;24(3):289-304
- Halsted CH et al (2002) Metabolic interactions of alcohol and folate. *J Nutr*. 2002 Aug;132(8 Suppl):2367S-2372S
- Halsted CH et al (2002b) Folate deficiency disturbs hepatic methionine metabolism and promotes liver injury in the ethanol-fed micropig. *Proc Natl Acad Sci U S A*. 2002 Jul 23;99(15):10072-7. Epub 2002 Jul 16
- Halsted CH et al (2002c) Interactions of ethanol and folate deficiency in development of alcoholic liver disease in the micropig. *Trans Am Clin Climatol Assoc*. 2002;113:151-62; discussion 162-3
- Halsted CH, Villanueva JA, Devlin AM (2002) Folate deficiency, methionine metabolism, and alcoholic liver disease. *Alcohol*. 2002 Jul;27(3):169-72
- Huang RF et al (2001) Folate depletion and elevated plasma homocysteine promote oxidative stress in rat livers. *J Nutr*. 2001 Jan;131(1):33-8
- James SJ et al (2003) Mechanisms of DNA damage, DNA hypomethylation, and tumor progression in the folate/methyl-deficient rat model of hepatocarcinogenesis. *J Nutr*. 2003 Nov;133(11 Suppl 1):3740S-3747S
- Ji C, Kaplowitz N (2004) Hyperhomocysteinemia, endoplasmic reticulum stress, and alcoholic liver injury. *World J Gastroenterol*. 2004 Jun 15;10(12):1699-708
- Joshi R et al (2001) Free radical scavenging behavior of folic acid: evidence for possible antioxidant activity. *Free Radic Biol Med*. 2001 Jun 15;30(12):1390-9
- Krishnaswamy K, Madhavan Nair K (2001) Importance of folate in human nutrition. *Br J Nutr*. 2001 May;85 Suppl 2:S115-24
- Niculescu MD, Zeisel SH (2002) Diet, methyl donors and DNA methylation: interactions between dietary folate, methionine and choline. *J Nutr*. 2002 Aug;132(8 Suppl):2333S-2335S
- O'Leary K, Sheehy PJ (2001) Influence of folic acid-fortified foods on folate status in a folate depletion-repletion rat model. *Br J Nutr*. 2001 Apr;85(4):441-6
- Poirier LA (2002) The effects of diet, genetics and chemicals on toxicity and aberrant DNA methylation: an introduction. *J Nutr*. 2002 Aug;132(8 Suppl):2336S-2339S
- Poschl G et al (2004) Alcohol and cancer: genetic and nutritional aspects. *Proc Nutr Soc*. 2004 Feb;63(1):65-71
- Poschl G, Seitz HK (2004) Alcohol and cancer. *Alcohol*. 2004 May-Jun;39(3):155-65
- Roncales M et al (2004) Folic acid supplementation for 4 weeks affects liver morphology in aged rats. *J Nutr*. 2004 May;134(5):1130-3
- Tapiero H et al (2001) Prevention of pathologies associated with oxidative stress and dietary intake deficiencies: folate deficiency and requirements. *Biomed Pharmacother*. 2001 Sep;55(7):381-90
- van Ede AE et al (2001) Effect of folic or folinic acid supplementation on the toxicity and efficacy of methotrexate in rheumatoid arthritis: a forty-eight week, multicenter, randomized, double-blind, placebo-controlled study. *Arthritis Rheum*. 2001 Jul;44(7):1515-24.
- Villanueva JA, Halsted CH (2004) Hepatic transmethylation reactions in micropigs with alcoholic liver disease. *Hepatology*. 2004 May;39(5):1303-10
- Walker SP, Wein P, Ihle BU (1997) Severe folate deficiency masquerading as the syndrome of hemolysis, elevated liver enzymes, and low platelets. *Obstet Gynecol*. 1997 Oct;90(4 Pt 2):655-7
- References for Vitamin B6**
- Baker H et al (1975) Inability of chronic alcoholics with liver disease to use food as a source of folates, thiamin and vitamin B6. *Am J Clin Nutr*. 1975 Dec;28(12):1377-80
- Benedikt J, Roth-Maier DA, Kirchgessner M (1996) The effect of different vitamin B6 supplies on the vitamin B6 status (pyridoxine, pyridoxal and pyridoxamine) of the liver and the body of lactating rats. *Z Ernährungswiss*. 1996 Sep;35(3):273-81
- Berrino F (2002) Western diet and Alzheimer's disease. *Epidemiol Prev*. 2002 May-Jun;26(3):107-15
- Bertrand J, Klos A, Debski B (2004) Influence of vitamin B6 supplementation on polyunsaturated fatty acids concentration in serum and liver of rats fed a diet restricted in protein. *Nahrung*. 2004 Apr;48(2):99-103
- Cabrini L et al (1998) Vitamin B6 deficiency affects antioxidant defences in rat liver and heart. *Biochem Mol Biol Int*. 1998 Nov;46(4):689-97
- Dastur DK, Dave UP (1987) Effect of prolonged anticonvulsant medication in epileptic patients: serum lipids, vitamins B6, B12, and folic acid, proteins, and fine structure of liver. *Epilepsia*. 1987 Mar-Apr;28(2):147-59
- Halsted CH (2004) Nutrition and alcoholic liver disease. *Semin Liver Dis*. 2004 Aug;24(3):289-304
- Hsu PC, Guo YL (2002) Antioxidant nutrients and lead toxicity. *Toxicology*. 2002 Oct 30;180(1):33-44
- Ji C, Kaplowitz N (2004) Hyperhomocysteinemia, endoplasmic reticulum stress, and alcoholic liver injury. *World J Gastroenterol*. 2004 Jun 15;10(12):1699-708
- Labadarios D et al (1977) Vitamin B6 deficiency in chronic liver disease—evidence for increased degradation of pyridoxal-5-phosphate. *Gut*. 1977 Jan;18(1):23-7
- Lau-Cam CA, Thadikonda KP, Kendall BF (1991) Stimulation of rat liver glycolysis by vitamin B6: a role for adrenal catecholamines. *Res Commun Chem Pathol Pharmacol*. 1991 Aug;73(2):197-207
- Meisler NT, Nutter LM, Thanassi JW (1982) Vitamin B6 metabolism in liver and liver-derived tumors. *Cancer Res*. 1982 Sep;42(9):3538-43
- Merrill AH Jr, Henderson JM (1990) Vitamin B6 metabolism by human liver. *Ann NY Acad Sci*. 1990;585:110-7
- Niculescu MD, Zeisel SH (2002) Diet, methyl donors and DNA methylation: interactions between dietary folate, methionine and choline. *J Nutr*. 2002 Aug;132(8 Suppl):2333S-2335S
- Oka T et al (1994) Effect of vitamin B6 deficiency on the expression of glycogen phosphorylase mRNA in rat liver and skeletal muscle. *Experientia*. 1994 Feb 15;50(2):127-9
- Okada M, Ishikawa K, Watanabe K (1991) Effect of vitamin B6 deficiency on glycogen metabolism in the skeletal muscle, heart, and liver of rats. *J Nutr Sci Vitaminol (Tokyo)*. 1991 Aug;37(4):349-57
- Poschl G, Seitz HK (2004) Alcohol and cancer. *Alcohol*. 2004 May-Jun;39(3):155-65
- Pregnotato P et al (1994) Interaction among dietary vitamin B6, proteins and lipids: effects on liver lipids in rats. *Int J Vitam Nutr Res*. 1994;64(4):263-9
- Roth-Maier DA et al (1996) Effect of feeding various dietary vitamin B6 concentrations during gestation and lactation on vitamin B6 level in liver and carcass of rat dams. *Arch Tierernahr*. 1996;49(4):319-24

References

References for Vitamin B12

Akesson B, Fehling C, Jagerstad M (1978) Effect of vitamin B12 deficiency on phosphatidylethanolamine methylation in rat liver. *Br J Nutr.* 1978 Nov;40(3):521-7

Areekul S, Panatampon P, Doungbarn J (1977) Vitamin B12 and vitamin B12 binding proteins in liver diseases. *Southeast Asian J Trop Med Public Health.* 1977 Sep;8(3):322-8

Baker H et al (1998) Cobalamin (vitamin B12) and holotranscobalamin changes in plasma and liver tissue in alcoholics with liver disease. *J Am Coll Nutr.* 1998 Jun;17(3):235-8

Berrino F (2002) Western diet and Alzheimer's disease. *Epidemiol Prev.* 2002 May-Jun;26(3):107-15

Brunaud L et al (2003) Effects of vitamin B12 and folate deficiencies on DNA methylation and carcinogenesis in rat liver. *Clin Chem Lab Med.* 2003 Aug;41(8):1012-9

Dastur DK, Dave UP (1987) Effect of prolonged anticonvulsant medication in epileptic patients: serum lipids, vitamins B6, B12, and folic acid, proteins, and fine structure of liver. *Epilepsia.* 1987 Mar-Apr;28(2):147-59

Fehling C et al (1978) Effects of vitamin B12 deficiency on lipid metabolism of the rat liver and nervous system. *Br J Nutr.* 1978 May;39(3):501-13

Ji C, Kaplowitz N (2004) Hyperhomocysteinemia, endoplasmic reticulum stress, and alcoholic liver injury. *World J Gastroenterol.* 2004 Jun 15;10(12):1699-708

Kennedy DG et al (1994) Cobalt-vitamin B12 deficiency causes lipid accumulation, lipid peroxidation and decreased alpha-tocopherol concentrations in the liver of sheep. *Int J Vitam Nutr Res.* 1994;64(4):270-6

McGing PG, Scott JM (1980) The role of methionine and vitamin B12 in folate incorporation by rat liver. *Br J Nutr.* 1980 Jan;43(1):235-7

Mikhail TH et al (1980) The effect of vitamin B12 on tetracycline-induced fatty liver. *Z Ernahrungswiss.* 1980 Sep;19(3):173-8

Niculescu MD, Zeisel SH (2002) Diet, methyl donors and DNA methylation: interactions between dietary folate, methionine and choline. *J Nutr.* 2002 Aug;132(8 Suppl):2333S-2335S

Okuda K (1999) Discovery of vitamin B12 in the liver and its absorption factor in the stomach: a historical review. *J Gastroenterol Hepatol.* 1999 Apr;14(4):301-8

Poirier LA (2002) The effects of diet, genetics and chemicals on toxicity and aberrant DNA methylation: an introduction. *J Nutr.* 2002 Aug;132(8 Suppl):2336S-2339S

Poznanskaia AA et al (1989) The monooxygenase system, lipid peroxidation in the liver and vitamin B12 metabolism in experimental poisoning of rabbits with phenylhydrazine. *Biokhimiia.* 1989 Aug;54(8):1290-3

Temcharoen P, Anukarahanonta T, Bhamarapravati N (1978) Influence of dietary protein and vitamin B12 on the toxicity and carcinogenicity of aflatoxins in rat liver. *Cancer Res.* 1978 Jul;38(7):2185-90

Toyoshima S et al (1996) Accumulation of methylmalonic acid caused by vitamin B12-deficiency disrupts normal cellular metabolism in rat liver. *Br J Nutr.* 1996 Jun;75(6):929-38

References for Ascorbic acid

Bsoul SA, Terezhalmay GT (2004) Vitamin C in health and disease. *J Contemp Dent Pract.* 2004 May 15;5(2):1-13

Carvalho M (2004) The toxicity of N-methyl-alpha-methyltyrosine to freshly isolated rat hepatocytes is prevented by ascorbic acid and N-acetylcysteine. *Toxicology.* 2004 Aug 5;200(2-3):193-203.

Celec P et al (2003) Effects of anabolic steroids and antioxidant vitamins on ethanol-induced tissue injury. *Life Sci.* 2003 Dec 12;74(4):419-34

Chopra M, Thurnham DI (1999) Antioxidants and lipoprotein metabolism. *Proc Nutr Soc.* 1999 Aug;58(3):663-71

Eguchi M et al (2003) Cytoprotection against ischemia-induced DNA cleavages and cell injuries in the rat liver by pro-vitamin C via hydrolytic conversion into ascorbate. *Mol Cell Biochem.* 2003 Oct;252(1-2):17-23

Elejalde Guerra JI (2001) Oxidative stress, diseases and

antioxidant treatment. *An Med Interna.* 2001 Jun;18(6):326-35

Farombi EO et al (2004) Commonly consumed and naturally occurring dietary substances affect biomarkers of oxidative stress and DNA damage in healthy rats. *Food Chem Toxicol.* 2004 Aug;42(8):1315-22

Flora SJ, Pande M, Mehta A (2003) Beneficial effect of combined administration of some naturally occurring antioxidants (vitamins) and thiol chelators in the treatment of chronic lead intoxication. *Chem Biol Interact.* 2003 Jun 15;145(3):267-80

George J (2003) Ascorbic acid concentrations in dimethylnitrosamine-induced hepatic fibrosis in rats. *Clin Chim Acta.* 2003 Sep;335(1-2):39-47

Hagymasi K, Blazovics A (2004) Antioxidants in liver protection. *Orv Hetil.* 2004 Jul 4;145(27):1421-5

Harrison SA et al (2003) Vitamin E and vitamin C treatment improves fibrosis in patients with nonalcoholic steatohepatitis. *Am J Gastroenterol.* 2003 Nov;98(11):2485-90

Hsu PC, Guo YL (2002) Antioxidant nutrients and lead toxicity. *Toxicology.* 2002 Oct 30;180(1):33-44

Kaneko K, Nagao N, Miwa N (1999) Vitamin C: structure-activity correlation and cytoprotective actions through free radical scavenging and extracellular matrix construction. *Nippon Rinsho.* 1999 Oct;57(10):2223-9

Kojo S (2004) Vitamin C: basic metabolism and its function as an index of oxidative stress. *Curr Med Chem.* 2004 Apr;11(8):1041-64

McDonough KH (2003) Antioxidant nutrients and alcohol. *Toxicology.* 2003 Jul 15;189(1-2):89-97

Moller P et al (2003) Oxidative DNA damage in vitamin C-supplemented guinea pigs after intratracheal instillation of diesel exhaust particles. *Toxicol Appl Pharmacol.* 2003 May 15;189(1):39-44

Muriel P, Moreno MG (2004) Effects of silymarin and vitamins E and C on liver damage induced by prolonged biliary obstruction in the rat. *Basic Clin Pharmacol Toxicol.* 2004 Feb;94(2):99-104

Patrick L (2003) Toxic metals and antioxidants: Part II. The role of antioxidants in arsenic and cadmium toxicity. *Altern Med Rev.* 2003 May;8(2):106-28

Ramanathan K et al (2003) Ascorbic acid and alpha-tocopherol as potent modulators on arsenic induced toxicity in mitochondria. *J Nutr Biochem.* 2003 Jul;14(7):416-20

Senthilkumar R, Viswanathan P, Nalini N (2004) Effect of glycine on oxidative stress in rats with alcohol induced liver injury. *Pharmazie.* 2004 Jan;59(1):55-60

Shugalei luS (1987) Features of vitamin C metabolism and the functional status of the liver in alcoholism and alcoholic delirium in the stage of detoxification therapy. *Zh Nevropatol Psikhiatr Im S S Korsakova.* 1987;87(2):240-3

Thirunavukkarasu V, Anuradha CV, Viswanathan P (2003) Protective effect of fenugreek (*Trigonella foenum graecum*) seeds in experimental ethanol toxicity. *Phytother Res.* 2003 Aug;17(7):737-43

Tselepidis S et al (2004) Liver injury after ischemia and reperfusion: the role of oxygen free radicals. *Mil Med.* 2004 Jul;169(7):531-5

Carvalho M (2004) The toxicity of N-methyl-alpha-methyltyrosine to freshly isolated rat hepatocytes is prevented by ascorbic acid and N-acetylcysteine. *Toxicology.* 2004 Aug 5;200(2-3):193-203.

Celec P et al (2003) Effects of anabolic steroids and antioxidant vitamins on ethanol-induced tissue injury. *Life Sci.* 2003 Dec 12;74(4):419-34

Eguchi M et al (2003) Cytoprotection against ischemia-induced DNA cleavages and cell injuries in the rat liver by pro-vitamin C via hydrolytic conversion into ascorbate. *Mol Cell Biochem.* 2003 Oct;252(1-2):17-23

Elejalde Guerra JI (2001) Oxidative stress, diseases and

References for Hesperidin

Cha JY et al (2001) Effect of hesperetin, a citrus flavonoid, on the liver triacylglycerol content and phosphatidate phosphohydrolase activity in orotic acid-fed rats. *Plant Foods Hum Nutr.* 2001;56(4):349-58

Chen YT et al (1990) Flavonoids as superoxide scavengers and antioxidants. *Free Radic Biol Med.* 1990;9(1):19-21

Chiba H et al (2003) Hesperidin, a citrus flavonoid, inhibits bone loss and decreases serum and hepatic lipids in ovariectomized mice. *J Nutr.* 2003 Jun;133(6):1892-7

Fraga CG et al (1987) Flavonoids as antioxidants evaluated by in vitro and in situ liver chemiluminescence. *Biochem Pharmacol.* 1987 Mar 1;36(5):717-20

Jung UJ et al (2004) The hypoglycemic effects of hesperidin and naringin are partly mediated by hepatic glucose-regulating enzymes in C57BL/KsJ-db/db mice. *J Nutr.* 2004 Oct;134(10):2499-503

Miyake Y et al (1998) Protective effects of lemon flavonoids on oxidative stress in diabetic rats. *Lipids.* 1998 Jul;33(7):689-95

Park YB et al (2001) Interactive effect of hesperidin and vitamin E supplements on cholesterol metabolism in high cholesterol-fed rats. *Int J Vitam Nutr Res.* 2001 Jan;71(1):36-44

References for Rutin

Afanas'ev IB et al (1995) Different antioxidant activities of bioflavonoid rutin in normal and iron-overloading rats. *Biochem Pharmacol.* 1995 Aug 25;50(5):627-35

Afanas'ev IB, Ostrachovich EA, Korkina LG (1998) Effect of rutin and its copper complex on superoxide formation and lipid peroxidation in rat liver microsomes. *FEBS Lett.* 1998 Mar 27;425(2):256-8

Aherne SA, O'Brien NM (1999) Protection by the flavonoids myricetin, quercetin, and rutin against hydrogen peroxide-induced DNA damage in Caco-2 and Hep G2 cells. *Nutr Cancer.* 1999;34(2):160-6

Bear WL, Teel RW (2000) Effects of citrus phytochemicals on liver and lung cytochrome P450 activity and on the in vitro metabolism of the tobacco-specific nitrosamine NNK. *Anticancer Res.* 2000 Sep-Oct;20(5A):3323-9

Boyle SP et al (2000) Bioavailability and efficiency of rutin as an antioxidant: a human supplementation study. *Eur J Clin Nutr.* 2000 Oct;54(10):774-82

Canada AT et al (1990) The production of reactive oxygen species by dietary flavonols. *Free Radic Biol Med.* 1990;9(5):441-9

Chen YT et al (1990b) Flavonoids as superoxide scavengers and antioxidants. *Free Radic Biol Med.* 1990;9(1):19-21

Fraga CG et al (1987b) Flavonoids as antioxidants evaluated by in vitro and in situ liver chemiluminescence. *Biochem Pharmacol.* 1987 Mar 1;36(5):717-20

Funabiki R et al (1999) Dietary supplement of G-rutin reduces oxidative damage in the rodent model. *J Agric Food Chem.* 1999 Mar;47(3):1078-82

Gao Z, Xu H, Chen X, Chen H (2003) Antioxidant status and mineral contents in tissues of rutin and baicalin fed rats. *Life Sci.* 2003 Aug 8;73(12):1599-607

Gao Z, Xu H, Huang K (2002) Effects of rutin supplementation on antioxidant status and iron, copper, and zinc contents in mouse liver and brain. *Biol Trace Elem Res.* 2002 Sep;88(3):271-9

Janbaz KH, Saeed SA, Gilani AH (2002) Protective effect of rutin on paracetamol- and CCl4-induced hepatotoxicity in rodents. *Fitoterapia.* 2002 Dec;73(7-8):557-63

Robak J, Gryglewski RJ (1988) Flavonoids are scavengers of superoxide anions. *Biochem Pharmacol.* 1988 Mar 1;37(5):837-41

Schwedhelm E et al (2003) Clinical pharmacokinetics of antioxidants and their impact on systemic oxidative stress. *Clin Pharmacokinet.* 2003;42(5):437-59

Siess MH, Vernevauf MF (1982) The influence of food flavonoids on the activity of some hepatic microsomal monooxygenases in rats. *Food Chem Toxicol.* 1982 Dec;20(6):883-6

van Acker FA et al (2000) Flavonoids can replace alpha-tocopherol as an antioxidant. *FEBS Lett.* 2000 May 12;473(2):145-8




Hepatic Detox

Love your liver and get more out of life!

Suggested Uses for Hepatic Detox:

- EXPOSURE TO HEAVY METALS & ENVIRONMENTAL TOXINS
- HIGH COPPER LEVELS
- CONSTIPATION
- LOOSE BOWELS
- INDIGESTION
- BLOATING
- POOR APPETITE
- FOOD ALLERGIES
- BAD BREATH
- PSORIASIS
- NAUSEA ASSOCIATED WITH FATTY MEALS
- FATTY LIVER
- CIRRHOSIS
- HEPATITIS
- HANGOVERS
- ALCOHOLICS
- BOWEL WIND
- HEADACHES / MIGRANES
- SUGAR CRAVINGS
- SKIN PROBLEMS
- DERMATITIS
- ACNE
- SKIN RASHES
- TIREDNESS
- FATIGUE
- INSOMNIA
- IMMUNE WEAKNESS
- DIET HIGH IN PROCESSED FOODS
- AGING PROCESS
- OBESITY
- WEIGHT LOSS
- STRESS
- LIGHT COLOURED STOOLS
- HYPERCHOLESTEROLEMIA
- DRY HAIR
- HAIR - SPLIT ENDS
- DRY SCALP
- GOUT
- JOINT PAINS
- SPOON NAILS
- NAILS SPLITTING OR LIFTING AT THE EDGES
- HORMONAL IMBALANCES
- PMT
- USERS OF ORAL CONTRACEPTIVE
- IRRITABILITY
- ANGER
- DEPRESSION
- ANXIETY
- DRUG USE e.g. PARACETAMOL
- BILE DISTURBANCES

Hepatic Detox is a comprehensive liver support nutrient required for good liver function and helps support detoxification.



Each tablet contains:

Silybum marianum ext. equiv Dry fruit (St. Mary's Thistle) Standardised to contain <i>Silybin</i>	3.5g 35mg	Magnesium Aspartate (Magnesium 1.7mg)	25mg
L. Methioine	100mg	Zinc Amino Acid Chelate (Zinc 4mg)	20mg
L. Glycine	75mg	Selenomethionine (selenium 5mcg)	12.5mg
L. Cysteine Hydrochloride	50mg	Folic Acid	25mcg
L. Glutamine	50mg	Pyridoxal-5-Phosphate	2.5mg
Inositol	50mg	Pyridoxine Hydrochloride (vitamin B6)	10mg
L. Lysine Hydrochloride	50mg	Cyanocobalamin (vitamin B12)	25mcg
L. Taurine	30mg	Ascorbic Acid (Vitamin C)	50mg
Levocarnitine	25mg	Hesperidin	40mg
Glutamic Acid	25mg	Rutin	10mg
Lecithin	40mg		
Potassium Aspartate (potassium 5.4mg)	25mg		

Proudly Distributed Throughout Australia:

N.S.W.	Page1 Wholesale Distributors	Phone 02 9736 1800	Fax 02 9736 3133
VIC. / TAS.	Oborne Health Supplies	Phone 03 8831 3888	Fax 03 8831 3898
S.A. / N.T.	Bettalife	Phone 08 8351 8455	Fax 08 8351 8722
QUEENSLAND	Fitness Products Natural Remedies Group	Phone 07 3849 8711 Phone 07 3889 8830	Fax 07 3849 8620 Fax 07 3889 8848
W.A.	David Woolcott	Phone 08 9385 6066	Fax 08 9385 6077